

TRIPLE LINKING NUMBERS AND HEEGAARD FLOER HOMOLOGY

EUGENE GORSKY, TYE LIDMAN, BEIBEI LIU, AND ALLISON H. MOORE

ABSTRACT. We establish some new relationships between Milnor invariants and Heegaard Floer homology. This includes a formula for the Milnor triple linking number from the link Floer complex, detection results for the Whitehead link and Borromean rings, and a structural property of the d -invariants of surgeries on certain algebraically split links.

1. INTRODUCTION

Milnor defined in his thesis a family of higher-order linking invariants capable of detecting triple linking, as in the Borromean rings [Mil54]. The μ -invariants are calculated using coefficients in the Magnus expansion of certain quotients of the fundamental group of the link complement. The Milnor invariants contain both link homotopy and concordance invariants of links [Mil57, Cas75, Sta65], and are central tools in the study of three-manifolds and four-manifolds. Geometric interpretations of Milnor's invariants are numerous. Stallings conjectured that the μ -invariants could be described in terms of Massey products for cohomology [Sta75], following which Turaev and Porter gave explicit interpretations [Tur76, Por80]. Milnor's invariants can be computed using the intersection theory of certain "derived" surfaces in the link exterior [Coc90a] or expressed in terms of the Alexander and Conway polynomials [Mur66, Tra84, Smy67]. The interpretation of the μ -invariants most relevant to our purposes is the identification of the first non-vanishing coefficient of the Conway polynomial of an algebraically split link with the Sato-Levine invariant β and square of the Milnor triple linking number μ_{123} for two- and three-component links, respectively, as determined by Cochran [Coc85].

As many other invariants from knot theory can be seen in Floer homology, it is natural to ask about the Milnor invariants as well. For example, it is asked in [OSS15, Problem 17.2.7]:

Problem 1.1. Do the Milnor invariants place algebraic restrictions on the structure of link Floer homology?

In previous work of the first, third, and fourth authors [GLM20], it is shown that Heegaard Floer homology is able to see the Sato-Levine invariant β of an algebraically split two-component link. In this paper, we address Problem 1.1 to study several appearances of the Milnor triple linking number μ_{123} [Mil57] in the Heegaard Floer theory of links and three-manifolds. We also apply this to give new Dehn surgery and link Floer complex detection results for the Whitehead link and the Borromean rings.

The first result we have is that the link Floer complex of Ozsváth-Szabó [OS08a] contains the Milnor triple linking number.

Theorem 1.2. *Let \mathcal{L} be a three-component link with pairwise linking number zero. Then, there is an explicit formula for $|\mu_{123}(\mathcal{L})|$ in terms of $CFL^-(\mathcal{L})$.*

For the explicit statement, see Corollary 3.10 below. The main strategy is to utilize the aforementioned relationship between the triple linking number and the Conway polynomial, and to express this in terms of the Euler characteristic of the link Floer complex and an associated link invariant.

If we add some additional hypotheses on the links then this formula simplifies greatly. The formula from Theorem 1.2 becomes especially transparent if the link is a Brunnian L -space link. First, recall that the h -function is an integer valued link invariant that is defined using absolute gradings in link Floer homology (see Section 2.2 for a definition and properties). Second, recall that a link of three or more components is called Brunnian if all its proper sublinks are unlinks. In this article, we will call a two-component link Brunnian if it is linking number zero and the components are unknots, and will generally include unlinks in the class of Brunnian links. More generally, a link is called algebraically split if all pairwise linking numbers are zero. Finally, a link is called an L -space link if all sufficiently large surgeries of S^3 are L -spaces, that is, they have the simplest possible Heegaard Floer homology. (In the first part of this section, we will work exclusively with \mathbb{Z}_2 -coefficients.)

Theorem 1.3. *Assume that \mathcal{L} is a Brunnian L -space link with three components. Then $\mu_{123}(\mathcal{L})^2 = \sum_{\mathbf{s}} h(\mathbf{s})$.*

In particular, $\sum_{\mathbf{s}} h(\mathbf{s})$ is a link homotopy invariant of three-component Brunnian L -space links. See Corollary 3.10 for a more general statement of homotopy invariance in the link Floer complex. For such links, the h -function is always non-negative (see Lemma 2.7 (7)). As the unlink is the only L -space link with vanishing h -function (Lemma 2.16), we have

Corollary 1.4. *Assume that \mathcal{L} is a Brunnian L -space link with three components. If $\mu_{123} = 0$ then \mathcal{L} is the three-component unlink.*

Next, we ask to what extent Floer homology is able to detect the simplest links admitting rationally framed Dehn surgery to the three-sphere or the Poincaré homology sphere. We first observe that if there is a rational surgery $S_{1/m_1, \dots, 1/m_\ell}^3(\mathcal{L})$ on an ℓ -component Brunnian link which is the three-sphere, then \mathcal{L} is the unlink (see Proposition 4.1). We extend this by considering rational surgeries which are the Poincaré homology sphere and prove the following results:

Theorem 1.5. *Let \mathcal{L} be an ℓ -component Brunnian link, and suppose that $S_{1/m_1, \dots, 1/m_\ell}^3(\mathcal{L})$ is the Poincaré homology sphere.*

- (1) *If $\ell = 2$, then \mathcal{L} is the Whitehead link or its mirror and $mn = 1$.*
- (2) *If $\ell = 3$, then \mathcal{L} is the Borromean rings, and all $|m_i| = 1$ with the same sign.*

(3) If $\ell \geq 4$, no such \mathcal{L} exists.

In a related vein, we prove that amongst Brunnian L-space links, the Whitehead link and Borromean rings are essentially detected by the Sato-Levine invariant and the triple linking number.

Theorem 1.6. *Let \mathcal{L} be an ℓ -component Brunnian L-space link.*

- (1) *If $\ell = 2$ and $\beta = 1$, then \mathcal{L} is the Whitehead link.*
- (2) *If $\ell = 3$ and $\mu_{123}(\mathcal{L}) = \pm 1$, then \mathcal{L} is the Borromean rings.*
- (3) *If $\ell \geq 4$, then \mathcal{L} is the four-component unlink.*

Finally, if we drop the assumption that the components of $\mathcal{L} = L_1 \cup L_2$ are unlinks, requiring only that it is an L-space link, we widen the detection result as follows.

Theorem 1.7. *Let $\mathcal{L} = L_1 \cup L_2$ be an algebraically split L-space link such that $S_{1,1}^3(\mathcal{L})$ is the Poincaré homology sphere. Then \mathcal{L} is either the Whitehead link or the split union of $T(2,3)$ and the unknot.*

We make the following conjecture for three-component algebraically split L-space links.

Conjecture 1.8. *Let $\mathcal{L} = L_1 \cup L_2 \cup L_3$ be an algebraically split L-space link such that $S_{1,1,1}^3(\mathcal{L})$ is the Poincaré homology sphere. Then \mathcal{L} must be one of the following:*

- (1) *the Borromean rings,*
- (2) *the split union of the Whitehead link and the unknot,*
- (3) *the split union of the right hand trefoil and the two-component unlink.*

Remark 1.9. By a similar argument to the one in [GLM20, Proposition 5.6], one of the manifolds $S_1^3(L_i)$ is an L-space where $i = 1, 2, 3$. Hence at least one of the components of L is the unknot or the right-handed trefoil $T(2,3)$.

Theorems 1.6 is sufficient to establish the following.

Corollary 1.10. *The link Floer chain complex detects the Whitehead link and the Borromean rings.*

Proof. The link Floer chain complex determines the Heegaard Floer homology of large surgeries on a link and hence determines whether a link is an L-space link. The link Floer chain complex also determines the multivariable Alexander polynomial [OS08a], which in turn determines the linking number [Tor53], as well as whether the link is Brunnian. By Theorem 1.6, it thus suffices to know that the link Floer complex determines the Sato-Levine invariant (for links with two components) or Milnor triple linking number (for links with three components). This is shown in [GLM20] for links with two components and Theorem 1.2 for links with three components. \square

The final appearance of the triple linking number involves its relationship with another Heegaard Floer-theoretic invariant, the d -invariant of a homology three-sphere. As is common, we write d for the d -invariant when computing Heegaard Floer homology with \mathbb{Z}_2 -coefficients. We will use $d_{\mathbb{k}}$ for the d -invariant when working with coefficients in a field \mathbb{k} . We show that the non-vanishing of the triple linking number for a link \mathcal{L} gives an interesting restriction on the d -invariants of integer homology spheres obtained by Dehn surgeries along \mathcal{L} in the three-sphere. We prove the following:

Theorem 1.11. *Let $\mathcal{L} = L_1 \cup L_2 \cup L_3$ be an algebraically split link such that all two-component sublinks are \mathbb{Q} -L-space links. If the triple linking number μ_{123} is nonzero, then $d_{\mathbb{Q}}(S_{1,1,1}^3(\mathcal{L})) \leq -2$. If the triple linking number μ_{123} is odd, then the analogous inequality holds with \mathbb{Z}_2 -coefficients.*

In Theorem 1.11, we use d -invariants for Heegaard Floer homology with coefficients in \mathbb{Q} and in \mathbb{Z}_2 . Although it is customary for Heegaard Floer homology literature to work over \mathbb{Z}_2 , we use a comparison with monopole Floer homology to obtain a separate result over \mathbb{Q} . Surprisingly enough, we do not know if $d_{\mathbb{Q}}$ and d coincide. However, note that a \mathbb{Z}_2 -L-space is necessarily a \mathbb{Q} -L-space. See Section 2.4 for a discussion about coefficients.

Remark 1.12. The same proof applies if we work with coefficients in an arbitrary field \mathbb{k} and the characteristic of \mathbb{k} is coprime to $\mu_{123}(\mathcal{L})$.

The L-space link assumption in Theorem 1.11 will be shown to be necessary in Example 5.21. As an immediate application of Theorem 1.11, notice that when $\mathcal{L} = L_1 \cup L_2 \cup L_3$ is an algebraically split link such that all two-component sublinks are L-space links and has non-vanishing triple linking number μ_{123} , then $S_{1,1,1}^3(\mathcal{L})$ has infinite order in homology cobordism group, and, for example, does not bound rational homology ball.

Remark 1.13. It is easy to see that Theorem 1.11 holds for algebraically split links \mathcal{L} with n components which contains a three-component sublink L_{ijk} satisfying the assumption of the theorem.

In Section 5 we establish some d -invariant inequalities for surgeries on links over an arbitrary field \mathbb{k} which may be of independent interest to the reader. Since link Floer homology is only defined over \mathbb{Z}_2 at the moment, for arbitrary coefficients, we cannot make use of it or various formulas relating link Floer homology with the Heegaard Floer homology of surgery. Nonetheless, we obtain the following results (which are known to experts over \mathbb{Z}_2):

Proposition 1.14. *Assume that \mathcal{L} is a nontrivial \mathbb{k} -L-space link of any number of components and pairwise linking zero. Then $d_{\mathbb{k}}(S_{1,\dots,1}^3(\mathcal{L})) \leq -2$. The same inequality is true for any $(1/m_1, \dots, 1/m_\ell)$ -surgery along \mathcal{L} where $m_1, \dots, m_\ell > 0$.*

Corollary 1.15. *Let \mathcal{L} be an algebraically split \mathbb{k} -L-space link such that $S_{1,\dots,1}^3(\mathcal{L})$ is S^3 . Then \mathcal{L} is the unlink.*

Organization. In Section 2, we review certain numerical invariants extracted from the link Floer complex, including (and defining) the h -function. In Section 3, we relate these numerical invariants to the Milnor invariants and prove Theorem 1.2. In Section 4 we establish the detection results for the Whitehead link and Borromean rings: Theorems 1.5, 1.6 and 1.7. Finally, in Section 5, we establish Theorem 1.11 on the d -invariants of surgeries on three-component algebraically split links.

Acknowledgements. We thank Marco Golla, Robert Lipshitz, Filip Misev, Yi Ni and Abigail Thompson for useful discussions. E.G. was supported by the grants DMS-1700814 and DMS-1760329. T.L. was supported by DMS-1709702 and a Sloan Fellowship. B.L. is grateful to Max Planck Institute for Mathematics in Bonn for its hospitality and financial support. A.H.M. was partially supported by DMS-1716987.

2. BACKGROUND

In this section, we review the relevant aspects of Heegaard Floer homology, especially properties of the link Floer complex and L-space links. A multi-component link is denoted by scripted \mathcal{L} , and its components are denoted L_i . We denote multi-framings and vectors in an n -dimensional lattice by bold letters (e.g. $\mathbf{p} = (p_1, \dots, p_n)$ -surgery or $\mathbf{s} = (s_1, \dots, s_n)$), and we let \mathbf{e}_i denote a vector in \mathbb{Z}^n where the i -th entry is 1 and other entries are 0. For any subset $B \subset \{1, \dots, n\}$, we let $\mathbf{e}_B = \sum_{i \in B} \mathbf{e}_i$. Given two vectors $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, \dots, v_n)$ in \mathbb{Z}^n , we write $\mathbf{u} \leq \mathbf{v}$ if $u_i \leq v_i$ for each $1 \leq i \leq n$, and $\mathbf{u} < \mathbf{v}$ if $\mathbf{u} \leq \mathbf{v}$ and $\mathbf{u} \neq \mathbf{v}$.

Definition 2.1. A link $\mathcal{L} = L_1 \cup \dots \cup L_n$ is algebraically split if for all $i \neq j$, $\ell k(L_i, L_j) = 0$.

Throughout this article, all links will be assumed to be algebraically split, unless otherwise stated. However, we include a slightly more general discussion below for the benefit of the reader.

Definition 2.2. For an oriented link $\mathcal{L} = L_1 \cup \dots \cup L_n \subset S^3$, define $\mathbb{H}(\mathcal{L})$ to be the affine lattice over \mathbb{Z}^n ,

$$\mathbb{H}(\mathcal{L}) = \bigoplus_{i=1}^n \mathbb{H}_i(\mathcal{L}), \quad \mathbb{H}_i(\mathcal{L}) = \mathbb{Z} + \frac{\ell k(L_i, \mathcal{L} \setminus L_i)}{2}.$$

If \mathcal{L} is algebraically split then $\mathbb{H}(\mathcal{L}) = \mathbb{Z}^n$.

2.1. d -invariants. We assume familiarity with Heegaard Floer homology, and refer the reader to [OS03, MO10] for details. With the exception of Section 5, we work over the field $\mathbb{F} = \mathbb{Z}_2$, as is typical in Heegaard Floer homology. Recall the d -invariant $d(Y, \mathfrak{t})$ of a rational homology sphere Y equipped with a Spin^c structure \mathfrak{t} is defined to be the maximal degree of a non-torsion class $x \in HF^-(Y, \mathfrak{t})$. For three-manifolds with $b_1(Y) > 0$, the definition of the d -invariant is more complicated, see Section 5.1.

2.2. The h -function and L -space links. We review the definition of the h -function for oriented links $\mathcal{L} \subset S^3$, as defined by the first author and Némethi [GN15]. We will quote without proof several technical lemmas regarding its properties; proofs of these statements can be found in either [BG18], [GLM20], or both.

A link $\mathcal{L} = L_1 \cup \cdots \cup L_n$ in S^3 defines a filtration on the Floer complex $CF^-(S^3)$. This filtration is indexed by elements $\mathbf{s} = (s_1, \dots, s_n) \in \mathbb{H}(\mathcal{L})$ (see Definition 2.2). The *generalized Heegaard Floer complex* $\mathfrak{A}^-(\mathcal{L}, \mathbf{s}) \subset CF^-(S^3)$ is the $\mathbb{F}[U]$ -module defined to be the subcomplex of $CF^-(S^3)$ corresponding to the filtration indexed by the lattice point \mathbf{s} [MO10]. The large surgery theorem of [MO10, Theorem 12.1] implies that the homology of $\mathfrak{A}^-(\mathcal{L}, \mathbf{s})$ is isomorphic to the Heegaard Floer homology of a sufficiently large surgery on the link \mathcal{L} equipped with some Spin^c -structure as an $\mathbb{F}[U]$ -module. Therefore, there is a non-canonical isomorphism between the homology of $\mathfrak{A}^-(\mathcal{L}, \mathbf{s})$ and a direct sum of one copy of $\mathbb{F}[U]$ and a U -torsion submodule. Thus the following is well-defined:

Definition 2.3. [BG18, Definition 3.9] For an oriented link $\mathcal{L} \subseteq S^3$, we define the H -function $H_{\mathcal{L}}(\mathbf{s})$ by saying that $-2H_{\mathcal{L}}(\mathbf{s})$ is the maximal homological degree of a nonzero element in the free part of $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{s}))$ where $\mathbf{s} \in \mathbb{H}(\mathcal{L})$.

Remark 2.4. We will write $H_{\mathcal{L}}(\mathbf{s})$ as $H(\mathbf{s})$ for brevity if the context is clear. The notation $H_I(\mathbf{s})$ refers to the H -function of the sublink determined by indices $I \subseteq \{1, \dots, n\}$.

By definition $-2H_{\mathcal{L}}(\mathbf{s})$ is equivalent to the d -invariant of large surgery on \mathcal{L} , following a degree shift that depends only on the linking matrix and \mathbf{s} (this is explained in [MO10, Section 10], [BG18, Theorem 4.10]). In particular, the H -function is a well-defined topological invariant of links. For a knot K , the H -function satisfies $H_K(\mathbf{s}) = V_{\mathbf{s}}$, where $V_{\mathbf{s}}$ are the similarly defined integer-valued concordance invariants coming from the knot Floer complex [Ras03, NW15].

We will work with a normalized version of the H -function, denoted h , as well as a ‘state sum’, denoted h' , that is taken over sublinks of \mathcal{L} .

Definition 2.5. Let \mathcal{L} be an n -component algebraically split link, $n \geq 1$. We define

$$\begin{aligned} h(\mathbf{s}) &= H(\mathbf{s}) - H_O(s_1) - \dots - H_O(s_n) \\ h'(\mathbf{s}) &= \sum_{I \subseteq \{1, \dots, n\}} (-1)^{n-|I|} h_I(\mathbf{s}) \end{aligned}$$

where $H_O(s)$ is the H -function for the unknot and $h(\emptyset) = 0$.

Example 2.6. Assume that \mathcal{L} is a Brunnian link, that is, all proper sublinks are unlinks. Then $h_I(\mathbf{s}) = 0$ for all proper subsets I and $h'(\mathbf{s}) = h(\mathbf{s})$.

We now list several properties of the H -function.

Lemma 2.7. For an oriented link $\mathcal{L} \subseteq S^3$,

- (1) The H -function $H_{\mathcal{L}}(\mathbf{s})$ takes nonnegative values.
- (2) $H_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_i) = H_{\mathcal{L}}(\mathbf{s})$ or $H_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_i) = H_{\mathcal{L}}(\mathbf{s}) + 1$ where $\mathbf{s} \in \mathbb{H}$.

(3) $H(-\mathbf{s}) = H(\mathbf{s}) + \sum_{i=1}^n s_i$ where $\mathbf{s} = (s_1, \dots, s_n)$.

(4) If \mathcal{L} is algebraically split and N is sufficiently large, then

$$H_{\mathcal{L}}(s_1, \dots, s_{i-1}, N, s_{i+1}, \dots, s_n) = H_{\mathcal{L} \setminus L_i}(s_1, \dots, s_{i-1}, s_{i+1}, \dots, s_n),$$

for all i and s_j .

(5) The functions h and h' have the following symmetry property:

$$h(-\mathbf{s}) = h(\mathbf{s}), \quad h'(-\mathbf{s}) = h'(\mathbf{s}).$$

(6) The function h is non-decreasing towards the origin. That is, $h(\mathbf{s} - \mathbf{e}_i) \geq h(\mathbf{s})$ if $s_i > 0$ and $h(\mathbf{s} - \mathbf{e}_i) \leq h(\mathbf{s})$ if $s_i \leq 0$.

(7) For all \mathbf{s} one has $h(\mathbf{s}) \geq 0$.

Proof. Items (1) and (2) are proved in [BG18, Proposition 3.10] and (3) is proved in [Liu17, Lemma 5.5]. Item (4) is [BG18, Proposition 3.12]. For (5), the first equation follows from (3) and the second follows from the first. Item (6) and Item (7) are proved in [GLM20, Lemma 2.16, Corollary 2.17]. \square

Lemma 2.8. *Let \mathcal{L} be an algebraically split link. Then the function $h'_{\mathcal{L}}(\mathbf{s})$ is finitely supported.*

Proof. By Lemma 2.7(4), when $s_i > N$ for some N , we have $h_{I \sqcup \{i\}}(\mathbf{s}) = h_I(\mathbf{s})$. By Lemma 2.7(5), it is therefore sufficient to prove for all i that when $s_i \rightarrow +\infty$ we have $h'(\mathbf{s}) = 0$. Fix any index i and observe that we can write

$$h'(\mathbf{s}) = \sum_{i \notin I} (-1)^{n-|I|} (h_I(\mathbf{s}) - h_{I \sqcup \{i\}}(\mathbf{s})). \quad \square$$

Lemma 2.9. *Suppose that \mathcal{L} is an n -component split link and $n > 1$. Then $h'(\mathbf{s}) = 0$ for all \mathbf{s} .*

Proof. Since \mathcal{L} is split, we have $h_I(\mathbf{s}) = \sum_{i \in I} h_i(s_i)$ for all I , and

$$h'(\mathbf{s}) = \sum_I (-1)^{n-|I|} \sum_{i \in I} h_i(s_i) = \sum_i h_i(s_i) \sum_{i \in I} (-1)^{n-|I|} = 0$$

for $n > 1$. \square

Corollary 2.10. *Let \mathcal{L} be an algebraically split link with $n > 1$ components. Then*

$$h'(s_1, \dots, s_n) = \sum_{I \subseteq \{1, \dots, n\}} (-1)^{n-|I|} H_I(\mathbf{s}).$$

Proof. If \mathcal{L} is an unlink, then similarly to Lemma 2.9 the right hand side vanishes, and then the statement follows by linearity. \square

Lemma 2.11. *Let \mathcal{L} be an algebraically split link with $n > 1$ components. Then for all \mathbf{s} one has*

$$\sum_{J \subseteq \{1, \dots, n\}} (-1)^{|J|} h'_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_J) = \sum_{J \subseteq \{1, \dots, n\}} (-1)^{|J|} H_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_J).$$

Proof. By Corollary 2.10 we get

$$\sum_{J \subset \{1, \dots, n\}} (-1)^{|J|} h'(\mathbf{s} - \mathbf{e}_J) = \sum_{I, J \subset \{1, \dots, n\}} (-1)^{n-|I|+|J|} H_I(\mathbf{s} - \mathbf{e}_J)$$

If I is a proper subset of $\{1, \dots, n\}$ then $H_I(\mathbf{s} - \mathbf{e}_J) = H_I(\mathbf{s} - \mathbf{e}_{I \cap J})$ and we can write $J = J' \sqcup J''$ where $J' = I \cap J$ and $J'' = J \setminus I$. Then

$$\sum_{J \subset \{1, \dots, n\}} (-1)^{n-|I|+|J|} H_I(\mathbf{s} - \mathbf{e}_J) = \sum_{J' \subset I} (-1)^{n-|I|+|J'|} H_I(\mathbf{s} - \mathbf{e}_{J'}) \sum_{J'' \subset (\{1, \dots, n\} \setminus I)} (-1)^{|J''|} = 0,$$

so the only surviving terms are from $I = \{1, \dots, n\}$. \square

2.3. L-spaces. Recall from [OS05] that a rational homology sphere Y is an L-space if it has the simplest possible Heegaard Floer homology. More precisely, for any Spin^c -structure \mathfrak{s} , $HF^-(Y, \mathfrak{s})$ is a free $\mathbb{F}[U]$ -module of rank one.

Definition 2.12. [GN15, Liu17] An oriented n -component link $\mathcal{L} \subset S^3$ is an L-space link if there exists $\mathbf{0} < \mathbf{p} \in \mathbb{Z}^n$ such that the surgered manifold $S_{\mathbf{q}}^3(\mathcal{L})$ is an L-space for any $\mathbf{q} \geq \mathbf{p}$.

Recall that if a knot $K \subset S^3$ admits any positive surgery to an L-space, then $S_{p/q}^3(K)$ is also an L-space for all $p/q \geq 2g(K) - 1$ [OS05]. For links though, it is not necessarily the case that the existence of a single \mathbf{p} -surgery yielding an L-space guarantees that all large surgeries are also L-spaces. However, the following criterion of Y. Liu can determine when this is the case.

Theorem 2.13. [Liu17]

- (1) *Every sublink of an L-space link is an L-space link.*
- (2) *A link is an L-space link if and only if for all \mathbf{s} one has $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{s})) = \mathbb{F}[U]$.*
- (3) *Assume that for some \mathbf{p} the surgery $S_{\mathbf{p}}^3(\mathcal{L})$ is an L-space. In addition, assume that for all sublinks $\mathcal{L}' \subset \mathcal{L}$ the surgeries $S_{\mathbf{p}|_{\mathcal{L}'}}^3(\mathcal{L}')$ are L-spaces too, and the framing matrix $\Lambda|_{\mathcal{L}'}$ is positive definite. Then for all $\mathbf{q} \geq \mathbf{p}$ the surgered manifolds $S_{\mathbf{q}}^3(\mathcal{L})$ are L-spaces, and so \mathcal{L} is an L-space link.*

Example 2.14. If \mathcal{L} is algebraically split, then Λ is positive definite if and only if all $p_i > 0$. In this case, the existence of any \mathbf{p} with all $p_i > 0$ such that $S_{\mathbf{p}}^3(\mathcal{L}')$ is an L-space for all sublinks \mathcal{L}' is sufficient to conclude \mathcal{L} is an L-space link. In particular, any Brunnian link admitting a single positive L-space surgery (e.g. $(+1, +1, +1)$ -surgery along the Borromean rings) is an L-space link.

By [OS08a], the Euler characteristic $\chi(HFL^-(\mathcal{L}, \mathbf{s}))$ is the multivariable Alexander polynomial,

$$(1) \quad \tilde{\Delta}_{\mathcal{L}}(t_1, \dots, t_n) = \sum_{\mathbf{s} \in \mathbb{H}(\mathcal{L})} \chi(HFL^-(\mathcal{L}, \mathbf{s})) t_1^{s_1} \cdots t_n^{s_n}$$

where $\mathbf{s} = (s_1, \dots, s_n)$, and

$$(2) \quad \tilde{\Delta}_{\mathcal{L}}(t_1, \dots, t_n) := \begin{cases} (t_1 \cdots t_n)^{1/2} \Delta_{\mathcal{L}}(t_1, \dots, t_n) & \text{if } n > 1, \\ \Delta_{\mathcal{L}}(t)/(1 - t^{-1}) & \text{if } n = 1. \end{cases}$$

For L-space links, the H -function can be computed from the multi-variable Alexander polynomial. Indeed, by Theorem 2.13 (2) and the inclusion-exclusion formula, one can write

$$(3) \quad \chi(HFL^-(\mathcal{L}, \mathbf{s})) = \sum_{B \subset \{1, \dots, n\}} (-1)^{|B|-1} H_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_B),$$

as in [BG18, (3.14)].

Example 2.15. The (symmetric) Alexander polynomial of the Whitehead link equals

$$\Delta(t_1, t_2) = -(t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2}),$$

and the H -function has the following values.

2	1	s_2 0	0	0
2	1	0	0	0
2	1	1	0	0 s_1
3	2	1	1	1
4	3	2	2	2

The H -function of the two-component unlink agrees everywhere with the H -function of the Whitehead link except at $\mathbf{s} = (0, 0)$, where $H_{\mathcal{O}}(\mathbf{0}) = 0$. Therefore, for the Whitehead link,

$$(4) \quad h_{\mathcal{L}}(s_1, s_2) = \begin{cases} 1 & \text{if } s_1 = s_2 = 0 \\ 0 & \text{otherwise.} \end{cases}$$

Lastly, we observe:

Lemma 2.16. *If for an L-space link \mathcal{L} one has $h(\mathbf{0}) = 0$, then \mathcal{L} is the unlink.*

Proof. If $h(\mathbf{0}) = 0$ then by Lemma 2.7 (6) we have $h(\mathbf{s}) = 0$ for all $\mathbf{s} \in \mathbb{H}(\mathcal{L})$. The rest of the proof follows from [Liu19a, Theorem 1.3]. \square

We will also make use of the following well-known fact without reference. If K is an L-space knot, then

$$g(K) = \max\{s \mid h(s) > 0\} + 1.$$

2.4. Coefficients. As stated above, for most of the paper we use $\mathbb{F} = \mathbb{Z}_2$ as the field of coefficients. However, in Section 5 we will use rational coefficients, so we need to discuss the dependence of the results on the field of coefficients.

First of all, the Heegaard Floer complexes $\widehat{CF}, CF^-, CF^\infty$ for knots and three-manifolds are defined over \mathbb{Z} [OS03]. In particular, CF^- is a complex of finitely generated free $\mathbb{Z}[U]$ modules. Its decomposition into Spin^c structures is well defined over $\mathbb{Z}[U]$.

Let \mathbb{k} be an arbitrary field. We will write $CF_{\mathbb{k}}^- = CF^- \otimes_{\mathbb{Z}} \mathbb{k}$, and define $\widehat{CF}_{\mathbb{k}}, CF_{\mathbb{k}}^\infty$ and $\widehat{HF}_{\mathbb{k}}, HF_{\mathbb{k}}^-, HF_{\mathbb{k}}^\infty$ similarly. Since $\mathbb{k}[U]$ is a principal ideal domain, any graded $\mathbb{k}[U]$ module (in particular, $HF_{\mathbb{k}}^-$) can be decomposed as a direct sum of several copies of $\mathbb{k}[U]$ and $\mathbb{k}[U]/U^{d_i}$ for various d_i .

If Y is a rational homology sphere, then for any Spin^c structure \mathfrak{t} on Y and any field \mathbb{k} the homology $HF_{\mathbb{k}}^-(Y, \mathfrak{t})$ contains exactly one copy of $\mathbb{k}[U]$ [OS04b]. We define $d_{\mathbb{k}}(Y, \mathfrak{t})$, the d -invariant with coefficients in \mathbb{k} , as the homological degree of the generator of this copy of $\mathbb{k}[U]$. When $\mathbb{k} = \mathbb{F} = \mathbb{Z}_2$, we simply write $d(Y, \mathfrak{t}) = d_{\mathbb{F}}(Y, \mathfrak{t})$, as above.

The following two examples show that d -invariants with coefficients in \mathbb{F} and in \mathbb{Q} could be potentially very different. It would be very interesting (but rather challenging) to find such examples in actual Heegaard Floer homology. In both examples we consider complexes of free $\mathbb{Z}[U]$ -modules with three generators a, b, c .

Example 2.17. Suppose that $\partial(c) = U^k a - 2b$. The homology over \mathbb{Z} can be identified with the submodule of $\mathbb{Z}[U]$ generated by 2 (corresponding to a) and U^k (corresponding to b). In particular, the homology is free as \mathbb{Z} -module and has no torsion.

On the other hand, if we consider this complex over \mathbb{F} , then $\partial(c) = U^k a$ and the homology is isomorphic to $\mathbb{F}[U]/(U^k) \oplus \mathbb{F}[U]$ as graded $\mathbb{F}[U]$ -module, its $\mathbb{F}[U]$ free part is generated by b . If we consider the same complex over \mathbb{Q} , then it is isomorphic to $\mathbb{Q}[U]$ generated by a .

In conclusion, $d_{\mathbb{F}} = d_{\mathbb{Q}} - 2k$.

Example 2.18. Suppose that $\partial(a) = U^k c$ and $\partial(b) = 2c$. In this case the homology has \mathbb{Z}_2 torsion of rank k , spanned by $c, Uc, \dots, U^{k-1}c$.

If we consider this complex over \mathbb{F} , then the homology is isomorphic to $\mathbb{F}[U]/(U^k) \oplus \mathbb{F}[U]$ as graded $\mathbb{F}[U]$ -module, and its $\mathbb{F}[U]$ free part is generated by b . If we consider the same complex over \mathbb{Q} , then it is isomorphic to $\mathbb{Q}[U]$ generated by $2a - U^k b$.

In conclusion, $d_{\mathbb{F}} = d_{\mathbb{Q}} + 2k$.

Note that the above examples show that the difference between $d_{\mathbb{F}}$ and $d_{\mathbb{Q}}$ could be either positive or negative, and arbitrarily large in absolute value.

It is also important to point out that that the notion of L-space (and hence of L-space link) depends on the coefficients, so a pedantically inclined reader is invited to use the terms \mathbb{F} -L-space and \mathbb{F} -L-space link.

At present, link Floer homology (for links with more than one component) is only defined over \mathbb{F} , so the H -function and its cousins are only defined over \mathbb{F} .

3. MILNOR INVARIANTS AND THE CASSON INVARIANT

In this section, we show how to extract the Milnor triple linking invariant from the link Floer complex. This will be in terms of the invariant h' defined in the previous section and another invariant χ' from the torsion part of $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{s}))$.

3.1. The invariant χ' . For non L-space links, the h -function does not determine the Alexander polynomial. However, we can obtain this from the collection of $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{s}))$ for all \mathbf{s} , which we now explain. Recall that for any link we have a non-canonical splitting

$$H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{s})) = \mathbb{F}[U][{-2H(\mathbf{s})}] \oplus \mathfrak{A}_{tor}^-(\mathcal{L}, \mathbf{s}),$$

where $\mathfrak{A}_{tor}^-(\mathcal{L}, \mathbf{s})$ is finite-dimensional over \mathbb{F} and hence a torsion module over $\mathbb{F}[U]$. We begin by analyzing the modules \mathfrak{A}_{tor}^- , as they will feature in our formula for the Alexander polynomial, and ultimately the Milnor invariants.

Lemma 3.1. *For an algebraically split link \mathcal{L} , we have*

$$H_*(\mathfrak{A}_{tor}^-(\mathcal{L}, -\mathbf{s})) \cong H_*(\mathfrak{A}_{tor}^-(\mathcal{L}, \mathbf{s}))[-2|\mathbf{s}|],$$

where $|\mathbf{s}| = \sum_i s_i$.

Proof. By the large surgery theorem [MO10] we have (up to a grading shift)

$$H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{s})) \cong HF^-(S_p^3(\mathcal{L}, \mathbf{s})), \text{ for } p \gg 0.$$

By [OS04b, Theorem 2.4] we have

$$HF^-(S_p^3(\mathcal{L}, -\mathbf{s})) \cong HF^-(S_p^3(\mathcal{L}, \mathbf{s}))$$

Therefore up to a grading shift we have

$$H_*(\mathfrak{A}^-(\mathcal{L}, -\mathbf{s})) \cong H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{s})).$$

To figure out the shift, we can look at the $\mathbb{F}[U]$ -free part and use the identity

$$H(-\mathbf{s}) = H(\mathbf{s}) + |\mathbf{s}|. \quad \square$$

Corollary 3.2. *For an algebraically split link \mathcal{L} , we have*

$$(5) \quad \chi(\mathfrak{A}_{tor}^-(\mathcal{L}, -\mathbf{s})) = \chi(\mathfrak{A}_{tor}^-(\mathcal{L}, \mathbf{s})).$$

We are ready to define the analogue of the function h' for the torsion parts of \mathfrak{A}^- .

Definition 3.3. Let \mathcal{L} be an algebraically split link with n components. We define

$$\chi'_{\mathcal{L}}(\mathbf{s}) = \chi'(\mathbf{s}) = \sum_{I \subset \{1, \dots, n\}} (-1)^{n-|I|} \chi(\mathfrak{A}_{tor}^-(\mathcal{L}_I, \mathbf{s}_I)).$$

Lemma 3.4. *The function $\chi'(\mathbf{s})$ is finitely supported and enjoys the symmetry $\chi'(-\mathbf{s}) = \chi'(\mathbf{s})$.*

Proof. The symmetry for $\chi'(\mathbf{s})$ immediately follows from (5). Let us prove that it is finitely supported. For $s_i \gg 0$ and any subset I not containing s_i we have from [MO10, Lemma 10.1]

$$H_*(\mathfrak{A}^-(\mathcal{L}_I, \mathbf{s}_I)) \cong H_*(\mathfrak{A}^-(\mathcal{L}_{I \cup \{i\}}, \mathbf{s}_{I \cup \{i\}}))$$

and, in particular,

$$\chi(\mathfrak{A}_{tor}^-(\mathbf{s}_I)) = \chi(\mathfrak{A}_{tor}^-(\mathbf{s}_{I \cup \{i\}})).$$

Then similarly to Lemma 2.8 we have $\chi'(\mathbf{s}) = 0$ for $s_i \gg 0$. By symmetry, we also have $\chi'(\mathbf{s}) = 0$ for $s_i \ll 0$, and therefore $\chi'(\mathbf{s})$ is finitely supported. \square

Finally, equation (3) can be generalized in the presence of torsion as follows:

$$(6) \quad \chi(HFL^-(\mathcal{L}, \mathbf{s})) = \sum_{B \subset \{1, \dots, n\}} (-1)^{|B|} (\chi(\mathfrak{A}_{tor}^-(\mathbf{s} - \mathbf{e}_B)) - H_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_B)).$$

Let \mathcal{L} be an algebraically split link with $n > 1$ components. Then the Torres condition [Tor53] implies that $\Delta_{\mathcal{L}}$ is divisible by $(t_i - 1)$ for all i . Hence, we can write

$$(7) \quad \Delta_{\mathcal{L}}(t_1, \dots, t_n) = \prod_i (t_i^{1/2} - t_i^{-1/2}) \tilde{\Delta}'_{\mathcal{L}}(t_1, \dots, t_n),$$

where $\Delta_{\mathcal{L}}$ is normalized as in equation (2) above.

Theorem 3.5. *Assume that \mathcal{L} is an algebraically split link with $n > 1$ components. Then*

$$\tilde{\Delta}'_{\mathcal{L}}(\mathbf{t}) = (-1)^n \sum_{\mathbf{s}} (\chi'(\mathbf{s}) - h'(\mathbf{s})) \mathbf{t}^{\mathbf{s}},$$

where $\mathbf{t}^{\mathbf{s}} = t_1^{s_1} \cdots t_n^{s_n}$.

Proof. Let $\tilde{\Delta}'_{\mathcal{L}}(\mathbf{t}) = \sum q(\mathbf{s}) \mathbf{t}^{\mathbf{s}}$ and $\tilde{\Delta}_{\mathcal{L}}(\mathbf{t}) = \prod_i (t_i - 1) \tilde{\Delta}'_{\mathcal{L}}(\mathbf{t}) = \sum a(\mathbf{s}) \mathbf{t}^{\mathbf{s}}$. Then

$$(8) \quad a(\mathbf{s}) = \sum_{J \subset \{1, \dots, n\}} (-1)^{n-|J|} q(\mathbf{s} - \mathbf{e}_J).$$

By Lemma 2.11 we get

$$\sum_{J \subset \{1, \dots, n\}} (-1)^{|J|} h'_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_J) = \sum_{J \subset \{1, \dots, n\}} (-1)^{|J|} H_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_J).$$

Similarly,

$$\sum_{J \subset \{1, \dots, n\}} (-1)^{|J|} \chi'(\mathbf{s} - \mathbf{e}_J) = \sum_{J \subset \{1, \dots, n\}} (-1)^{|J|} \chi(\mathfrak{A}_{tor}^-(\mathbf{s} - \mathbf{e}_J)).$$

Therefore (6) implies

$$(9) \quad a(\mathbf{s}) = \sum_{J \subset \{1, \dots, n\}} (-1)^{|J|} (\chi'(\mathbf{s} - \mathbf{e}_J) - h'(\mathbf{s} - \mathbf{e}_J)).$$

Equations (8) and (9) imply that $(-1)^n q(\mathbf{s})$ and $\chi'(\mathbf{s}) - h'(\mathbf{s})$ satisfy the same recursion relations, and by Lemmas 2.8 and 3.4 both vanish for sufficiently large \mathbf{s} . Therefore,

$$\tilde{\Delta}'_{\mathcal{L}}(\mathbf{t}) = (-1)^n \sum_{\mathbf{s}} (\chi'(\mathbf{s}) - h'(\mathbf{s})) \mathbf{t}^{\mathbf{s}}. \quad \square$$

Corollary 3.6. *Assume that \mathcal{L} is a Brunnian L -space link with $n > 1$ components. Then*

$$\tilde{\Delta}'_{\mathcal{L}}(\mathbf{t}) = (-1)^{n+1} \sum_{\mathbf{s}} h(\mathbf{s}) \mathbf{t}^{\mathbf{s}}.$$

Proof. For Brunnian links $h'(\mathbf{s}) = h(\mathbf{s})$, and since \mathcal{L} is an L -space link, $\chi'(\mathbf{s}) = 0$ for all \mathbf{s} . \square

3.2. Milnor triple linking invariant. To associate the Milnor triple linking invariant to the Alexander polynomial, we must pass through the Conway polynomial. The Conway polynomial of $\mathcal{L} = L_1 \cup \dots \cup L_n$ can be written as

$$\nabla_{\mathcal{L}}(z) = z^{n-1}(a_0 + a_2 z^2 + a_4 z^4 + \dots), \quad a_i \in \mathbb{Z},$$

and

$$\nabla_{\mathcal{L}}(t^{1/2} - t^{-1/2}) = \pm(t^{1/2} - t^{-1/2})\Delta_{\mathcal{L}}(t, \dots, t),$$

where $\Delta_{\mathcal{L}}(t_1, \dots, t_n)$ denotes the multi-variable Alexander polynomial of \mathcal{L} . Note that a_0 depends only on the linking numbers of \mathcal{L} , see [Hos85, Theorem 1]. For an algebraically split link with $n > 1$ components, $a_0 = 0$ and we can write its multivariable Alexander polynomial as in (7):

$$\Delta_{\mathcal{L}}(t_1, \dots, t_n) = \prod_i (t_i^{1/2} - t_i^{-1/2}) \tilde{\Delta}'_{\mathcal{L}}(t_1, \dots, t_n).$$

Then we can set all $t_i = t$ and apply the change of variable $z = t^{1/2} - t^{-1/2}$:

$$\nabla_{\mathcal{L}}(z) = \pm(t^{1/2} - t^{-1/2})\Delta_{\mathcal{L}}(t, \dots, t) = \pm(t^{1/2} - t^{-1/2})^{n+1} \tilde{\Delta}'_{\mathcal{L}}(t, \dots, t).$$

We define $\tilde{\nabla}_{\mathcal{L}}(z)$ as $\nabla_{\mathcal{L}}(z)/z^{n+1}$. With this, the coefficient a_2 of the Conway polynomial can be written as

$$a_2(\mathcal{L}) = \tilde{\nabla}_{\mathcal{L}}(0) = \pm \tilde{\Delta}'_{\mathcal{L}}(1, \dots, 1).$$

It is an important invariant of the link. By Theorem 3.5 we get

$$(10) \quad a_2(\mathcal{L}) = \pm \sum_{\mathbf{s}} (\chi'(\mathbf{s}) - h'(\mathbf{s})).$$

Example 3.7. For two-component algebraically split links the invariant $a_2(\mathcal{L})$ agrees with the Sato-Levine invariant $\beta(\mathcal{L})$ up to sign [GLM20, Stu84]. Equation (10) now gives an explicit formula for $\beta(\mathcal{L})$ in terms of the link Floer complex for \mathcal{L} . Moreover, if \mathcal{L} is a two-component algebraically split L -space link with unknotted components, then $\beta(\mathcal{L}) = 0$ implies \mathcal{L} is the unlink (see [GLM20, Corollary 6.4]).

Remark 3.8. For two-component links, other Milnor invariants of the form $\mu[1^p 2^q]$ may be written in terms of the link Floer complex as follows. (For example, the linking number corresponds with $\mu[12]$ and the Sato-Levine invariant with $\mu[1122]$.) One first writes the multivariable Alexander polynomial as in Theorem 3.5. Then passing to the Taylor expansion at $(1, 1)$ of this two variable polynomial, a result of Murasugi [Mur66, Theorem 4.1] shows that the coefficients of the Taylor expansion determine these Milnor invariants.

Example 3.9. If \mathcal{L} is an algebraically split link with $n \geq 4$ components then by [Coc85] $a_2(\mathcal{L}) = 0$.

Proof of Theorem 1.2. For three-component links, by [Coc85, Theorem 5.1] $a_2(\mathcal{L})$ relates to the square of the Milnor triple linking number

$$(11) \quad a_2(\mathcal{L}) = \pm \mu_{123}^2(\mathcal{L}).$$

Equation (10) now establishes an explicit formula for $|\mu_{123}|$ in terms of the link Floer complex. \square

Corollary 3.10. *Assume that \mathcal{L} is an algebraically split link with three components. Then*

$$\mu_{123}^2(\mathcal{L}) = \pm \sum_{\mathbf{s}} (\chi'(\mathbf{s}) - h'(\mathbf{s})).$$

In particular, $|\sum_{\mathbf{s}} (\chi'(\mathbf{s}) - h'(\mathbf{s}))|$ is a link homotopy invariant.

For Brunnian L -space links we can get even more information.

Theorem 3.11. *Assume that \mathcal{L} is a Brunnian L -space link with three components. Then the following statements hold:*

- (a) $\mu_{123}^2(\mathcal{L}) = \sum_{\mathbf{s}} h(\mathbf{s})$.
- (b) If $\mu_{123}(\mathcal{L}) = 0$ then \mathcal{L} is the unlink.
- (c) $\mu_{123}(\mathcal{L})$ has the same parity as $h(0, 0, 0) = H(0, 0, 0)$.
- (d) $\mu_{123}(\mathcal{L})$ cannot equal ± 2 .

Proof. (a) By (11) we have $\mu_{123}^2(\mathcal{L}) = \pm a_2(\mathcal{L})$ and by Corollary 3.10 we have $a_2(\mathcal{L}) = \sum_{\mathbf{s}} h(\mathbf{s})$. Therefore

$$\mu_{123}^2(\mathcal{L}) = \pm \sum_{\mathbf{s}} h(\mathbf{s}).$$

On the other hand, both μ_{123}^2 and $h(\mathbf{s})$ are nonnegative, so the sign is positive.

(b) If $\mu_{123}(\mathcal{L}) = 0$ then by (a) we have $h(\mathbf{s}) = 0$ for all \mathbf{s} . By [Liu19a, Theorem 1.3] this implies that \mathcal{L} is the unlink.

(c) By Lemma 2.7(5), we have $h(-\mathbf{s}) = h(\mathbf{s})$ for all \mathbf{s} . Note that $\mathbf{s} = -\mathbf{s}$ if and only if $\mathbf{s} = (0, 0, 0)$. Therefore $\sum_{\mathbf{s}} h(\mathbf{s})$ has the same parity as $h(0, 0, 0)$.

(d) Assume that $\mu_{123}(\mathcal{L}) = 2$, then by (c) $h(0, 0, 0)$ is even as well. If $h(0, 0, 0) = 0$ then $h(\mathbf{s}) = 0$ for all \mathbf{s} by Lemma 2.7(6), which is a contradiction. Therefore $h(0, 0, 0) \geq 2$, hence

$$h(1, 0, 0), h(0, 1, 0), h(0, 0, 1) \geq 1$$

by Lemma 2.7. This implies $\mu_{123}^2 = \sum_{\mathbf{s}} h(\mathbf{s}) \geq 2+3 = 5$, contradicting to the assumption. \square

Note that Theorem 3.11(a) and (b) provide the statement of Theorem 1.3 and Corollary 1.4 in the introduction. We mention here an open problem suggested by the discussion above:

Problem 3.12. Do there exist examples of Brunnian L -space links with three components and large μ_{123} ?

Let $\mathcal{L} = L_1 \cup L_2 \cdots \cup L_n$ be an oriented link in an integer homology sphere Y with all pairwise linking numbers equal zero, and with framing $1/q_i$ on component L_i , for $q_i \in \mathbb{Z}$. Hoste [Hos86] proved that the Casson invariant λ of the integer homology sphere $Y_{1/q_1, \dots, 1/q_n}(\mathcal{L})$ satisfies a state sum formula,

$$(12) \quad \lambda(Y_{1/q_1, \dots, 1/q_n}(\mathcal{L})) = \lambda(Y) + \sum_{\mathcal{L}' \subset \mathcal{L}} \left(\prod_{i \in \mathcal{L}'} q_i \right) a_2(\mathcal{L}'; Y),$$

where the sum is taken over all sublinks \mathcal{L}' of \mathcal{L} .

For example, let $\mathcal{L} = L_1 \cup L_2 \cup L_3$ be a three-component algebraically split link in S^3 with framings $q_i = 1$. Formula (12) simplifies to

$$\lambda(S_{1,1,1}^3(\mathcal{L})) = a_2(\mathcal{L}) + a_2(L_{12}) + a_2(L_{13}) + a_2(L_{23}) + a_2(L_1) + a_2(L_2) + a_2(L_3),$$

where $L_{ij} = L_i \cup L_j$. Theorem 3.11(a) immediately implies

Corollary 3.13. *Assume that \mathcal{L} is a Brunnian L -space link with three components. Then*

$$\lambda(S_{1,1,1}^3(\mathcal{L})) = \mu_{123}^2(\mathcal{L}) = \sum_{\mathbf{s}} h_{\mathcal{L}}(\mathbf{s}).$$

Example 3.14. Let \mathcal{L} denote the Borromean rings, which are easily checked to form an L -space link. Since $S_{1,1,1}^3(\mathcal{L})$ is the Poincaré homology sphere, we see that $\lambda(S_{1,1,1}^3(\mathcal{L})) = 1$. This confirms the well-known calculation that $|\mu_{123}(\mathcal{L})| = 1$. From this, we also deduce that the h -function satisfies $h(0, 0, 0) = 1$ and vanishes elsewhere.

4. DETECTION RESULTS

In this section, we will apply the statements from the section above to show that for Brunnian links or algebraically split L -space links, sometimes information about surgery or Milnor invariants is sufficient for link detection. In Section 4.1, we focus on Whitehead link characterizations, in Section 4.2, we focus on Borromean ring detections, and finally, in Section 4.3, we discuss analogous results for links with more components. Combining

the results in these sections, we obtain proofs of Theorem 1.5, Theorem 1.6, and Theorem 1.7.

Before we move to these statements, as a warm-up, we first determine which Brunnian links admit a rational surgery which is the three-sphere S^3 .

Proposition 4.1. *Let $\mathcal{L} = L_1 \cup \cdots \cup L_\ell$ be an ℓ -component Brunnian link, and suppose that some surgery $S_{1/m_1, \dots, 1/m_\ell}^3(\mathcal{L})$ is the three sphere S^3 , where all $m_i \neq 0$. Then \mathcal{L} is the unlink.*

Proof. We first prove that $S_{1, \dots, 1}^3(\mathcal{L}) = S^3$. Let L'_ℓ be the image of L_ℓ in $S_{1/m_1, \dots, 1/m_{\ell-1}}^3$ which is S^3 since \mathcal{L} is Brunnian. Then L'_ℓ is the unknot by Gordon-Luecke [GL89]. It is easy to see that

$$S_{1/m_1, \dots, 1/m_{\ell-1}, 1}^3(\mathcal{L}) = S_1^3(L'_\ell) = S^3.$$

By repeating this argument, one can easily prove that $S_{1, \dots, 1}^3(\mathcal{L}) = S^3$. By Example 2.14, \mathcal{L} is an L-space link. Hence \mathcal{L} is the unlink by Corollary 1.15. \square

In the following sections, we will generalize to the case when $(1/m_1, \dots, 1/m_\ell)$ -framed Dehn surgery yields the Poincaré homology sphere. We will use the notation $\text{PHS} = \Sigma(2, 3, 5)$ for the Poincaré homology sphere, oriented as the boundary of the positive-definite $E8$ plumbing, and Wh and B for the positive Whitehead and Borromean links, respectively.

4.1. Whitehead link detection. This subsection is devoted to the detection of the Whitehead link. Proposition 4.2 informs and precedes the more general Proposition 4.5, which gives the statement of Theorem 1.5(1) in the introduction. Corollary 4.7 will give the statement of Theorem 1.6(1), and following this we prove Theorem 1.7.

Proposition 4.2. *Let \mathcal{L} be a two-component Brunnian link. If $S_{1,1}^3(\mathcal{L})$ is the Poincaré homology sphere, then \mathcal{L} is the positive Whitehead link Wh .*

Proof. First, since \mathcal{L} is Brunnian, each of the components L_1 and L_2 are unknotted. Therefore, $S_n^3(L_i) = L(n, 1)$ for all integers n . For notation, let $E^\mathcal{L}$ denote the exterior of \mathcal{L} , and let T_1 and T_2 denote the boundary components corresponding to L_1 and L_2 respectively. Let $L_{2,(n)}$ denote the image of L_2 in $S_n^3(L_1)$ and $M_{(n)}^\mathcal{L}$ the exterior of $L_{2,(n)}$ in $S_n^3(L_1)$. Note that $M_{(n)}^\mathcal{L}$ has one torus boundary component, obtained by Dehn filling $E^\mathcal{L}$ along T_1 . We also will study $M_{(\infty)}^\mathcal{L}$, which is just a solid torus, since $L_{2,(\infty)}$ is unknotted. Since $S_{1,1}^3(\mathcal{L})$ is the Poincaré homology sphere, $L_{2,(1)}$ is the trefoil by [Ghi08]. Therefore, $M_{(1)}^\mathcal{L}$ is the exterior of the trefoil in S^3 , and hence is a fibered three-manifold.

We first claim that for all $n \neq \infty$, $L_{2,(n)}$ is a genus one fibered knot. From the above discussion, we see that there are two slopes α, β on the boundary component T_1 in $E^\mathcal{L}$ such that Dehn filling along α results in a fibered three-manifold and filling along β lowers the Thurston norm. (Here, with respect to the canonical meridian-longitude coordinates on T_1 , $\alpha = +1$ and $\beta = \infty$.) By [Ni09, Theorem 1.4], we see that the core J of the α -filling sits on the fiber surface F of the trefoil in $M_{(1)}^\mathcal{L}$ and that the framing

β corresponds to the surface framing of F . It follows that any surgery on J which is distance one from β is also a genus one fibered three-manifold. Indeed, one cuts along F and reglues by some number of Dehn twists along J . In meridian-longitude coordinates, J is the core of $+1$ -surgery on L_1 in the exterior of L_2 , β corresponds to the ∞ -filling of T_1 , and integral n -filling along L_1 for any n is distance one from β . Therefore each $M_{(n)}^{\mathcal{L}}$ is a fibered three-manifold with fiber having genus one and a single boundary component. Since $L_{2,(n)}$ is nullhomologous, we see that it is a genus one fibered knot in $L(n, 1)$.

By [Bak14, Theorem 4.3], for $n \neq 4, \infty$, we see that $L_{2,(n)}$ belongs to one of exactly two isotopy classes of knots. These must be either $\text{Wh}_{2,(n)}$ or $\overline{\text{Wh}}_{2,(n)}$ because the arguments above imply that integral surgery on a single component of Wh and $\overline{\text{Wh}}$ give genus one fibered knots. To see these are distinct, note that $+1$ -surgery on $\text{Wh}_{2,(n)}$ gives $S_n^3(T_{2,3})$ whereas $+1$ -surgery on $\overline{\text{Wh}}_{2,(n)}$ gives $S_n^3(4_1)$.

In other words, $M_{(n)}^{\mathcal{L}} = M_{(n)}^{\text{Wh}}$ or $M_{(n)}^{\overline{\text{Wh}}}$ for infinitely many n . By Lemma 4.3 below, $E^{\mathcal{L}} = E^{\text{Wh}}$ or $E^{\overline{\text{Wh}}}$. In principle, this does not yet imply that \mathcal{L} is a Whitehead link, since such links are not determined by their exteriors. However, Lemma 4.4 shows that the additional condition that $S_{1,1}^3(\mathcal{L})$ being the Poincaré homology sphere implies \mathcal{L} is in fact Wh . \square

Lemma 4.3. *If $M_{(n)}^{\mathcal{L}} = M_{(n)}^{\text{Wh}}$ for infinitely many n , then $E^{\mathcal{L}} = E^{\text{Wh}}$ and similarly for $\overline{\text{Wh}}$.*

Proof. We do the case of Wh . The mirror is the same.

Suppose $E^{\mathcal{L}}$ is hyperbolic with two cusps. Then by Thurston's hyperbolic Dehn surgery theorem [Thu97], $M_{(n)}^{\mathcal{L}}$ is hyperbolic for all but finitely many n with one cusp. In addition, $E^{\mathcal{L}}$ is the geometric limit of the sequence of complete hyperbolic manifolds $M_{(n)}^{\mathcal{L}}$ which is the same as $M_{(n)}^{\text{Wh}}$. Hence, in the limit $E^{\mathcal{L}} = E^{\text{Wh}}$.

We next prove that $E^{\mathcal{L}}$ is hyperbolic. Suppose instead that $E^{\mathcal{L}}$ is not hyperbolic. If $E^{\mathcal{L}}$ is reducible, then \mathcal{L} is split, and so \mathcal{L} is an unlink, and so $M_{(n)}^{\mathcal{L}} \neq M_{(n)}^{\text{Wh}}$, contradiction. Note that $E^{\mathcal{L}}$ is not Seifert, since otherwise $M_{(n)}^{\mathcal{L}}$ would not be hyperbolic for any n . It follows that there is a non-boundary parallel incompressible torus (i.e. essential torus) in $E^{\mathcal{L}}$. Consider the piece X of the JSJ decomposition of $E^{\mathcal{L}}$ which contains T_1 . Let T' denote a boundary component of X which is not T_1 or T_2 and essential in $E^{\mathcal{L}}$. Then, T' remains incompressible and non-boundary parallel in all but finitely many fillings of X along T_1 , unless X is a cable space (i.e. a Seifert fibered space with base orbifold an annulus and exactly one cone point) [CGLS87, Theorem 2.4.4]. If X is not a cable space, then generically $M_{(n)}^{\mathcal{L}}$ has a non-boundary parallel incompressible torus. However, $M_{(n)}^{\text{Wh}}$ is hyperbolic for infinitely many n , and hence does not have a non-boundary parallel incompressible torus, contradiction. If instead X is a cable space, and infinitely many of the n -fillings of T_1 are not those that cause T' to compress in $M_{(n)}^{\mathcal{L}}$, then again $M_{(n)}^{\mathcal{L}}$ has a non-boundary parallel incompressible torus for infinitely many n , while $M_{(n)}^{\text{Wh}}$ does not, contradiction.




FIGURE 1. The generalized twist knot $K[2n, -2m]$ is a two-bridge knot described by numerator closure of the rational tangle with continued fraction $[2n, -2m]$. For example, $K[2, -2]$ is the right-handed trefoil.

Lastly, suppose that X is a cable space, and that infinitely many integral n -slopes on L_1 are slopes on X which cause T' to compress. The fiber slope on a boundary component of a cable space is uniquely characterized by the slope which is distance one from at least 3 different compressing slopes [CGLS87, Theorem 2.4.3]. This means that the ∞ -slope on L_1 corresponds to the fiber slope ϕ on X . Recall that for a cable space, filling one boundary component along the fiber slope results in the connected sum of a solid torus and a non-trivial lens space. Let $X(\phi)$ denotes the Dehn filling of T_1 with the fiber slope ϕ . Therefore, $M_{(\infty)}^{\mathcal{L}}$ contains $X(\phi)$ as a codimension zero submanifold which has a lens space summand, but $M_{(\infty)}^{\mathcal{L}}$ is a solid torus, so this is a contradiction. \square

Lemma 4.4. *If \mathcal{L} is a two-component link in S^3 with $E^{\mathcal{L}} = E^{\text{Wh}}$ or $E^{\overline{\text{Wh}}}$ and $S_{+1,+1}^3(\mathcal{L})$ is the Poincaré homology sphere, then $\mathcal{L} = \text{Wh}$.*

To explain why this lemma is necessary, notice that doing $1/n$ -surgery on a single component of Wh yields S^3 , and that the image of the other component is a twist knot $K[2n, -2]$. The core of that surgery, together with this twist knot, is a two-component link in S^3 with the same exterior as Wh .

Proof. We begin with the case that $E^{\mathcal{L}} = E^{\text{Wh}}$. Since Wh has linking number 0, any \mathcal{L} in S^3 with the same exterior can be described by the core of $(1/m, 1/n)$ -surgery on Wh , where that surgery results in S^3 . Note that $S_{\frac{1}{m}, \frac{1}{n}}^3(\text{Wh})$ is $1/m$ -surgery on the twist knot $K[2n, -2]$ (see Figure 1), and thus either $1/m = \infty$ or $1/n = \infty$. Without loss of generality, $1/n = \infty$. Therefore,

$$\text{PHS} = S_{1,1}^3(\mathcal{L}) = S_{\frac{1}{m+1}, 1}^3(\text{Wh}) = S_{\frac{1}{m+1}}^3(K[2, -2]) = S_1^3(K[2(m+1), -2]).$$

It follows that $K[2(m+1), -2]$ is the right-handed trefoil, and hence $m = 0$.

We can repeat a similar argument with $\overline{\text{Wh}}$. In this case, we see that

$$\text{PHS} = S_{1,1}^3(\mathcal{L}) = S_{\frac{1}{m+1}, 1}^3(\overline{\text{Wh}}) = S_{\frac{1}{m+1}}^3(4_1) = S_1^3(4_1),$$

which is a contradiction. \square

By an argument similar to the one in Proposition 4.2, we now can characterize which rational surgeries on a Brunnian two-component link are the Poincaré homology sphere.

Proposition 4.5. *Suppose that \mathcal{L} is a two-component Brunnian link. Then $S_{1/m,1/n}^3(\mathcal{L})$ is the Poincaré homology sphere if and only if $mn = 1$ and \mathcal{L} is the Whitehead link or its mirror.*

Proof. The if part is easy. For the only if part, we first claim that if $S_{1/m,1/n}^3(\mathcal{L})$ is the Poincaré homology sphere, then $m = \pm 1$ and $n = \pm 1$. Suppose $\mathcal{L} = L_1 \cup L_2$, and let L'_1 denote the image of L_1 in S^3 after blowing down L_2 . Then $S_{1/m,1/n}^3(\mathcal{L}) = S_{1/m}^3(L'_1) = \pm\text{PHS}$, which implies that $m = \pm 1$, since only a trefoil has a surgery to the Poincaré homology sphere, and the surgery coefficient is integral [Ghi08]. A similar argument applies for n . If $m = n = 1$, by Proposition 4.2, \mathcal{L} must be the Whitehead link. If $m = n = -1$, we have $S_{-1,-1}^3(\mathcal{L}) = S_{1,1}^3(\bar{\mathcal{L}})$ where $\bar{\mathcal{L}}$ is the mirror of \mathcal{L} . Then $\bar{\mathcal{L}}$ must be the Whitehead link, and hence, \mathcal{L} is the mirror of the Whitehead link.

Without loss of generality, we assume that $m = 1, n = -1$. By a similar argument as the one in Proposition 4.2, the exterior of the link $E^{\mathcal{L}} = E^{Wh}$ or $E^{\overline{Wh}}$. However, this is impossible by applying the same argument as in Lemma 4.4. \square

We now transition to characterizing the Whitehead link among two-component algebraically split L-space links.

Lemma 4.6. *Let $\mathcal{L} = L_1 \cup L_2$ be an algebraically split L-space link with the following h -function:*

$$(13) \quad h_{\mathcal{L}}(s_1, s_2) = \begin{cases} 1 & \text{if } s_1 = s_2 = 0 \\ 0 & \text{otherwise.} \end{cases}$$

Then \mathcal{L} is the Whitehead link.

Proof. Let us prove that $S_{1,1}^3(\mathcal{L})$ is the Poincaré homology sphere. Let L'_2 be the image of L_2 in $S^3(L_1)$, so that $S_{1,1}^3(\mathcal{L}) = S_{1,1}^3(L'_2)$.

Since the h -function of \mathcal{L} is given by (13), the link surgery complex for $S_{1,1}^3(\mathcal{L})$ can be truncated so that it contains only $\mathfrak{A}_{00}^-(\mathcal{L})$ [GLM20]. Therefore $S_{1,1}^3(L'_2) = S_{1,1}^3(\mathcal{L})$ is an L-space with d -invariant equal to $-2H(0,0) = -2$. The knot L'_2 is an L-space knot and the L-space surgery coefficient of $+1$ satisfies $1 \geq 2g(L'_2) - 1$. Because the surgery $S_{1,1}^3(L'_2)$ is not S^3 , we must have the genus of L'_2 is exactly 1, and so L'_2 is the right-handed trefoil. This also shows $S_{1,1}^3(\mathcal{L})$ is the Poincaré homology sphere. Finally, since the unknot is determined by its h -function among L-space knots, Lemma 2.7(4) implies that the components of \mathcal{L} are unknotted. Now the statement follows from Proposition 4.2. \square

Corollary 4.7. *Suppose that \mathcal{L} is a two-component L-space link with unknotted components and $\beta(\mathcal{L}) = \pm 1$. Then \mathcal{L} is the Whitehead link.*

Proof. Since \mathcal{L} is an L-space link with unknotted components, we have $\sum_{\mathbf{s}} h(\mathbf{s}) = \pm\beta(\mathcal{L}) = \pm 1$. Since $h(\mathbf{s}) \geq 0$ and $h(-\mathbf{s}) = h(\mathbf{s})$, the only possible h -function is given by (13). Hence, \mathcal{L} is the Whitehead link by Lemma 4.6. \square

Theorem 1.7. *Let $\mathcal{L} = L_1 \cup L_2$ be an algebraically split L-space link such that $S_{1,1}^3(\mathcal{L})$ is the Poincaré homology sphere. Then \mathcal{L} is either the Whitehead link or the split union of $T(2, 3)$ and the unknot.*

Proof. By [GLM20, Proposition 5.6] we have that either $S_1^3(L_1)$ or $S_1^3(L_2)$ is an L-space. Without loss of generality, in the remainder of the proof we assume that $S_1^3(L_1)$ is an L-space. Then L_1 is an L-space knot of genus 0 or 1, so it is either unknotted or the right-handed trefoil.

Case 1: L_1 is an unknot. In this case we can blow it down and obtain a knot L'_2 such that $S_{1,1}^3(\mathcal{L}) = S_1^3(L'_2)$. This means that $S_1^3(L'_2)$ is the Poincaré homology sphere. So L'_2 is $T(2, 3)$.

By [GLM20, Theorem 4.8] the H -function for L'_2 equals $H_{\mathcal{L}}(0, s_2)$, so

$$h_{\mathcal{L}}(0, s_2) = h_{T(2,3)}(s_2) = \begin{cases} 1 & \text{if } s_2 = 0 \\ 0 & \text{otherwise.} \end{cases}$$

By Lemma 2.7(6) we get $h_{\mathcal{L}}(N, s_2) = 0$ for all $N \geq 0$ and $s_2 \neq 0$, and hence $h_2(s_2) = 0$ for $s_2 \neq 0$. Since L_2 is an L-space knot, this implies the genus is at most one, and hence it is either unknotted or $T(2, 3)$.




FIGURE 2. The h -function for $O \sqcup T(2, 3)$

If L_2 is unknotted, then \mathcal{L} is the Whitehead link by Proposition 4.2.

If L_2 is $T(2, 3)$, then $h_2(0) = h_{\mathcal{L}}(N, 0) = 1$ for all $N > 0$, and Lemma 2.7(5) implies that $h_{\mathcal{L}}(s_1, s_2)$ coincides with the h -function of the disjoint union of $O \sqcup T(2, 3)$ (see

Figure 2). By the main result of [Liu19b] the Thurston polytope of \mathcal{L} is the same as the Thurston polytope of $O \sqcup T(2, 3)$. In particular, L_1 bounds a disk not intersecting L_2 , and hence $\mathcal{L} = O \sqcup T(2, 3)$.

Case 2: L_1 is the trefoil. Let L'_2 be the knot corresponding to L_2 in $S^3_1(L_1) = \text{PHS}$. Since $+1$ -framed surgery along $L'_2 \subset \text{PHS}$ yields PHS , by the Dehn surgery characterization of the unknot in an integer homology sphere L-space [KMOS07, Gai18], L'_2 is an unknot.

Now for all d we have $S^3_{1,d}(\mathcal{L}) = \text{PHS}_d(L'_2) = S^3_{1,d}(T(2, 3) \sqcup O)$. Similarly to [GLM20, Theorem 4.8] we conclude that

$$h_{\mathcal{L}}(0, s_2) = h_{T(2,3) \sqcup O}(0, s_2) = 1$$

as in Figure 3.

Now we claim that L_2 is the unknot. Otherwise, suppose that $g(L_2) \geq 1$. By Lemma 2.7(6), $h_2(s_2) = h_{\mathcal{L}}(N, s_2) \leq h_{\mathcal{L}}(0, s_2) = 1$. More precisely, $h_2(s_2) = 0$ for all $|s_2| \geq g(L_2)$ and $h_2(s_2) = 1$ otherwise as in Figure 3.

Recall that $h'(s_1, s_2) = h_{\mathcal{L}}(s_1, s_2) - h_1(s_1) - h_2(s_2)$. Since L_1 is the trefoil, $h_1(s_1) = 0$ for all $s_1 \neq 0$. By a similar argument to the one in Lemma 2.7(6), one can prove that $h'(\mathbf{s} - \mathbf{e}_1) \geq h'(\mathbf{s})$ for all $s_1 > 1$. Note that $h'(\infty, s_2) = 0$ for all $s_2 \in \mathbb{Z}$. Then $h'(s_1, s_2) \geq 0$ for all $s_1 \geq 1$. By Lemma 2.7(5), $h'(s_1, s_2) \geq 0$ for all $|s_1| \geq 1$.

By (12),

$$1 = \lambda(S^3_{1,1}(\mathcal{L})) = a_2(\mathcal{L}) + a_2(L_1) + a_2(L_2).$$

Note that $a_2(L_1) = \sum h_1(s_1) = 1$, $a_2(L_2) = \sum h_2(s_2) = 2g(L_2) - 1$. Then $a_2(\mathcal{L}) = \sum h'_{\mathcal{L}}(\mathbf{s}) = 1 - 2g(L_2) < 0$. Observe that (see Figure 3)

$$\sum_{s_2 \in \mathbb{Z}} h'_{\mathcal{L}}(0, s_2) = 1 - 2g(L_2) = \sum_{\mathbf{s} \in \mathbb{H}(\mathcal{L})} h'_{\mathcal{L}}(s_1, s_2),$$

and $h'_{\mathcal{L}}(s_1, s_2) \geq 0$ for all $|s_1| \geq 1$. Hence $h'_{\mathcal{L}}(s_1, s_2) = 0$ for all $s_1 \neq 0$, indicating that $h_{\mathcal{L}}(s_1, s_2) = h_1(s_1) + h_2(s_2)$ for all $s_1 \neq 0$. Note that $h_1(s_1) = 0$ for all $s_1 \neq 0$. This implies that $h_{\mathcal{L}}(s_1, s_2) = h_2(s_2)$ for all $s_1 \geq 1$, and the h -function has the form as in Figure 3.

As in [GLM20, Section 5] we may define

$$b_1 = \min\{s_1 - 1 \mid H_{\mathcal{L}}(s_1, s_2) = H_2(s_2) \text{ for all } s_2\}.$$

Clearly, from Figure 3 we have $b_1 = 0$. It is proved in [Liu19c, Proposition 4.7] that (under some assumptions on the h -function which are satisfied in this case) if $S^3_{d_1, d_2}(\mathcal{L})$ is an L-space for $d_1 > 2b_1$ and $d_2 \ll 0$ then L_2 is the unknot. Since $(1, d)$ -surgery on \mathcal{L} yields $\text{PHS} \# L(d, 1)$, which is an L-space for any nonzero integer d , L_2 is the unknot. Hence, by the same argument as the one in Case 1, \mathcal{L} is the disjoint union of the unknot and $T(2, 3)$. \square

$$\begin{array}{ccccccc}
& & & \uparrow & & & \\
& 0 & \cdots & 0 & 1 & 0 & \cdots & 0 \\
& \vdots & & \vdots & \vdots & \vdots & & \vdots \\
& 0 & & 0 & 1 & 0 & & 0 \\
& 1 & & 1 & 1 & 1 & & 1 \leftarrow \cdots g(L_2) - 1 \\
\cdots & \vdots & & \vdots & \vdots & \vdots & \cdots & \vdots \\
& 1 & & 1 & 1 & 1 & & 1 \leftarrow \cdots 1 - g(L_2) \\
& 0 & & 0 & 1 & 0 & & 0 \\
& \vdots & & \vdots & \vdots & \vdots & & \vdots \\
& 0 & \cdots & 0 & 1 & 0 & \cdots & 0
\end{array}$$

FIGURE 3. Hypothetical h -function for Case 2

4.2. Borromean link detection. In this subsection, Proposition 4.8 will inform the more general Proposition 4.10 which corresponds with Theorem 1.5(2) in the introduction. Proposition 4.11 will give the statement of Theorem 1.6(2).

Proposition 4.8. *Let \mathcal{L} be a three-component Brunnian link. Then if $S_{1,1,1}^3(\mathcal{L})$ is the Poincaré homology sphere, then \mathcal{L} is the Borromean rings B .*

Proof. Because $L_1 \cup L_2 \cup L_3$ is Brunnian, all proper two-component sublinks $L_i \cup L_j$ are unlinks and $S_{m,n}^3(L_i \cup L_j) = L(m,1) \# L(n,1)$ for all integers m, n . We will use notation similar to that of Proposition 4.2. Let $E^{\mathcal{L}}$ denote the exterior of \mathcal{L} , and T_i the boundary component corresponding to L_i . We write $L_{3,(m),(n)}$ to denote the image of L_3 in $S_{m,n}^3(L_1 \cup L_2) = L(m,1) \# L(n,1)$, and write $M_{(n),(m)}^{\mathcal{L}}$ for the exterior of $L_{3,(m),(n)}$ in $L(m,1) \# L(n,1)$. We also write $L_{3,2,(n)}$ to denote the image of $L_2 \cup L_3$ in $S_n^3(L_1)$. We similarly have that $M_{(\infty),(\infty)}^{\mathcal{L}}$ is a solid torus. Because $S_{1,1,1}^3(\mathcal{L})$ is the Poincaré homology sphere, $L_{3,(1),(1)}$ is the trefoil knot and $M_{(1),(1)}^{\mathcal{L}}$ is the exterior of the trefoil in S^3 , a fibered three-manifold.

We will invoke [Ni09, Theorem 1.4] to argue that for all pairs of integers $n, m \neq \infty$, $L_{3,(n),(m)}$ is a genus one fibered knot. The argument is analogous to that of Proposition 4.2. There are two slopes α_1, β_1 on the boundary component T_1 in $E^{\mathcal{L}}$ such that Dehn filling along α_1 results in a fibered three-manifold and filling along β_1 lowers the Thurston norm. With respect to the canonical meridian-longitude coordinates on T_1 , $\alpha_1 = +1$ and $\beta_1 = \infty$. Indeed, filling along α_1 results in the complement of Whitehead link in S^3 . Since \mathcal{L} is Brunnian, then $L_{3,2,(1)}$ is also Brunnian and $S_{1,1}^3(L_{3,2,(1)}) = S_{1,1,1}^3(\mathcal{L})$ is the Poincaré homology sphere. By Proposition 4.2, $L_{3,2,(1)}$ is the Whitehead link. Filling along β_1 results in the complement of two-component unlink in S^3 , which lowers the Thurston norm. By [Ni09, Theorem 1.4], the core J_1 of the α_1 -filling of T_1 sits on the

fiber surface F of the Whitehead link complement $M_{(1)}^{\mathcal{L}}$. The framing β_1 corresponds to the surface framing of F . Now we repeat this argument. In particular, $M_{(1)}^{\mathcal{L}}$ has two boundary components T_2 and T_3 . We consider two fillings α_2 and β_2 along T_2 similarly (i.e. $\alpha_2 = +1, \beta_2 = \infty$). By the argument given above, the α_2 filling $M_{(1),(1)}^{\mathcal{L}}$ yields a fibered three-manifold which is the exterior of the right-handed trefoil and is obtained by capping off one boundary of the fiber surface of the Whitehead link. The β_2 filling $M_{(\infty),(1)}^{\mathcal{L}}$ which results in the complement of the unknot in S^3 reduces the Thurston norm. By a second application of [Ni09, Theorem 1.4], we have that the core J_2 of the α_2 -filling of T_2 sits on the fiber surface F_3 in $M_{(1),(1)}^{\mathcal{L}}$ and any surgery which is distance one from β_2 will produce a fibered three-manifold. In fact, any fillings along T_1, T_2 which are distance one from β_1, β_2 will extend the monodromy of the fibration. We now have for any integers m, n that $M_{(m),(n)}^{\mathcal{L}}$ is fibered and that the fiber is genus one. So $L_{3,(m),(n)}$ is a genus one fibered knot in $L(m, 1) \# L(n, 1)$.

Having established that $L_{3,(m),(n)}$ is genus one fibered knot, we turn again to Baker's classification of genus one fibered knots in lens spaces [Bak14]. By [Bak14, Lemma 2.2], the isotopy classes (M, K) of genus one fibered knots K in M are in a one-to-one correspondence with isotopy classes (N, A) where N is a link in S^3 whose branched double cover is M , and with a braid axis A that yields a closed 3-braid presentation of N . Fix m, n odd and distinct. Applying this here, we must have that N is the connected sum of torus knots $T(2, m) \# T(2, n)$. Indeed, $T(2, m)$ is the unique knot whose branched double cover is the lens space $L(m, 1)$ by Hodgson and Rubinstein [HR85]. By the equivariant sphere theorem for involutions [KT80], $T(2, m) \# T(2, n)$ is the unique knot with branched double cover $L(m, 1) \# L(n, 1)$. Thus, we seek to identify the braid axes A yielding closed 3-braid presentations for $T(2, m) \# T(2, n)$. There is a unique braid axis for closed 3-braid presentations of the knot $T(2, m) \# T(2, n)$ with odd integers satisfying $|m|, |n| > 1$. Note that the torus knots $T(2, m)$ and $T(2, n)$ are strongly invertible, so $T(2, m) \# T(2, n)$ is invertible. There is no distinction between the orientations of the knot. By [Bak14, Lemma 3.6], there is at most one equivalence class of braid axes giving 3-braid representatives for the oriented knot and its inverse.

Similarly to the argument used for the Whitehead link, we have that $M_{(m),(n)}^{\mathcal{L}} = M_{(m),(n)}^{\text{B}}$ for infinitely many m, n . An argument completely analogous to that of Lemma 4.3 implies that the geometric limit $E^{\mathcal{L}}$ of the hyperbolic manifolds $M_{(m),(n)}^{\mathcal{L}}$ agrees with the limit E^{B} of the sequence $M_{(m),(n)}^{\text{B}}$, meaning the exterior $E^{\mathcal{L}}$ is the same as the Borromean exterior E^{B} . The result now follows from Lemma 4.9 below. \square

Lemma 4.9. *If \mathcal{L} is a three-component Brunnian link in S^3 with $E^{\mathcal{L}} = E^{\text{B}}$ and $S_{1,1,1}^3(\mathcal{L})$ is the Poincaré homology sphere, then $\mathcal{L} = \text{B}$.*

Proof. Any link \mathcal{L} in S^3 with the same exterior as B can be described by the core of $(1/m, 1/n, 1/p)$ -surgery on B , where this surgery results in S^3 . By Proposition 4.1, at least one of m, n, p is zero. Without loss of generality, we assume $p = 0$. Then

$$\text{PHS} = S_{1,1,1}^3(\mathcal{L}) = S_{\frac{1}{m+1}, \frac{1}{n+1}, 1}^3(\text{B}) = S_{\frac{1}{m+1}, \frac{1}{n+1}}^3(\text{Wh}).$$

By Proposition 4.5, $m = n = 0$. Therefore, $\mathcal{L} = \mathbf{B}$. \square

Based on the argument for the detection of the Borromean rings, we now characterize which rational surgeries on a Brunnian three-component link are the Poincaré homology sphere.

Proposition 4.10. *Suppose that \mathcal{L} is a three-component Brunnian link. Then $S_{1/p, 1/q, 1/r}^3(\mathcal{L})$ is the Poincaré homology sphere if and only if the link \mathcal{L} is the Borromean rings, and $|p| = |q| = |r| = 1$ with the same sign.*

Proof. The if part is easy. For the only if part, we first claim that $p, q, r = \pm 1$. Suppose that $\mathcal{L} = L_1 \cup L_2 \cup L_3$, and let L'_1 denote the image of L_1 in S^3 after blowing down L_2, L_3 . Then $S_{1/p, 1/q, 1/r}^3(\mathcal{L}) = S_{1/p}^3(L'_1)$, which implies that $1/p = \pm 1$. Hence, $p = \pm 1$. Further, the sign of p is determined by the definiteness of the plumbing which the surgered Poincaré homology sphere bounds. A similar argument can be used to prove that $q, r = \pm 1$, and that their signs must agree with that of p . If $p = q = r = 1$, by Proposition 4.8, \mathcal{L} is the Borromean rings. If $p = q = r = -1$, then the mirror of \mathcal{L} is the Borromean rings, which implies that \mathcal{L} is also the Borromean rings. \square

Proposition 4.11. *Assume that \mathcal{L} is a Brunnian L -space link with three components and $\mu_{123}(\mathcal{L}) = \pm 1$. Then \mathcal{L} is the Borromean rings.*

Proof. By Theorem 3.11(a), $\sum_{\mathbf{s}} h(\mathbf{s}) = 1$. Recall that $h(\mathbf{s}) \geq 0$, and takes the maximal value at $h(0, 0, 0)$. So

$$h(\mathbf{s}) = \begin{cases} 1 & \text{if } \mathbf{s} = (0, 0, 0) \\ 0 & \text{otherwise.} \end{cases}$$

Let us prove that $S_{1,1,1}^3(\mathcal{L})$ is the Poincaré homology sphere. Because \mathcal{L} is Brunnian and the h -function agrees with that of the Borromean rings, the link surgery complex can be truncated so that it contains only $\mathfrak{A}_{000}^-(\mathcal{L})$. Just as with the proof of Lemma 4.6, because $S_{1,1,1}^3(\mathcal{L})$ is an L -space with d -invariant $-2H(0, 0, 0) = -2$ we again have that $S_{1,1,1}^3(\mathcal{L})$ is the Poincaré sphere. That is, by doing $+1$ -surgery along the link components L_1 and L_2 , we have a knot $L_3'' \subset S^3$ with a $+1$ -surgery to an L -space with $d = -2$. Thus L_3'' is the trefoil and $S_{1,1,1}^3(\mathcal{L})$ is the Poincaré sphere. By Proposition 4.8, \mathcal{L} is isotopic to the Borromean rings. \square

4.3. Links with four or more components. In this section we show that the above results have no analogues for links with more than three components. Proposition 4.12 gives Theorem 1.6(3) and Proposition 4.13 gives Theorem 1.5(3) in the introduction.

Proposition 4.12. *Assume that \mathcal{L} is a Brunnian L -space link with four or more components. Then \mathcal{L} is the unlink.*

Proof. By Example 3.9 we have $a_2(\mathcal{L}) = \tilde{\Delta}'(1, \dots, 1) = 0$. Therefore by Corollary 3.6 we have $\sum_{\mathbf{s}} h(\mathbf{s}) = 0$ and $h(\mathbf{s})$ vanishes for all \mathbf{s} . Then by [Liu19a, Theorem 1.3] \mathcal{L} is the unlink. \square

Now we consider whether rational surgery on a Brunnian link with at least four components is the Poincaré homology sphere. Note that if \mathcal{L} is a Brunnian link and $S_{1,\dots,1}^3(\mathcal{L})$ is an L-space, then by Theorem 2.13 it is an L-space link. By Proposition 4.12, \mathcal{L} is the unlink, and $(1, \dots, 1)$ -surgery cannot be the Poincaré homology sphere. For general rational surgeries, we have the similar result.

Proposition 4.13. *Let \mathcal{L} be an n -component Brunnian link with $n \geq 4$. Then $S_{1/m_1, \dots, 1/m_n}^3(\mathcal{L})$ is not the Poincaré homology sphere.*

Proof. Suppose $S_{1/m_1, \dots, 1/m_n}^3(\mathcal{L})$ is the Poincaré homology sphere. By the same argument as in Proposition 4.10, $m_i = \pm 1$ where $i = 1, \dots, n$. Since $S_{1/m_1, \dots, 1/m_n}^3(\mathcal{L}) = S_{1/m_2, \dots, 1/m_n}^3(L'_2 \cup \dots \cup L'_n)$ and the latter is still a Brunnian link, it suffices to consider the case that \mathcal{L} has four components. By repeating the arguments and notation from Proposition 4.2 and Proposition 4.8, we see that $L_{4,(l),(p),(q)}$ is a genus 1 fibered knot in (l, p, q) -surgery on the three-component unlink whenever l, p, q are integers. However, $L(l, 1) \# L(p, 1) \# L(q, 1)$ does not contain a genus 1 fibered knot since its Heegaard genus is 3 when $|l|, |p|, |q| > 1$, which is a contradiction. Hence, $S_{1/m_1, \dots, 1/m_n}^3(\mathcal{L})$ is not the Poincaré homology sphere. \square

5. FROM TRIPLE LINKING TO d -INVARIANTS

In this section, unlike the rest of the paper, we will use d -invariants for Heegaard Floer homology with a variety of coefficients. See Section 2.4 for more details on subtleties of coefficient fields. Some of the results hold over an arbitrary field \mathbb{k} , and we will denote the corresponding d -invariants by $d_{\mathbb{k}}$.

Let us also introduce some additional notation required in this section. Given a null-homologous link $\mathcal{L} = L_1 \cup \dots \cup L_\ell$ in a rational homology sphere Y and integers $\mathbf{m} = (m_1, \dots, m_\ell)$, let $W_{\mathbf{m}}$ denote the four-dimensional two-handle cobordism from Y to $Y_{\mathbf{m}}(\mathcal{L})$ which is the surgery manifold specified by the \mathbf{m} -framed link \mathcal{L} . The notation $\mathfrak{s}, \mathfrak{t}$ and $\mathfrak{w}, \mathfrak{z}$ will generally be used to denote Spin^c -structures on three-manifolds and four-manifolds, respectively. In particular, let \mathfrak{w} denote a Spin^c -structure on $W_{\mathbf{m}} = W_{\mathbf{m}}(K)$ which extends \mathfrak{s} in $\text{Spin}^c(Y)$, where K is a knot in Y . Recall from [OS08b, Theorem 4.2] that \mathfrak{w} induces a Spin^c -structure \mathfrak{s}_i on $Y_m(K)$ indexed by

$$\langle c_1(\mathfrak{w}), [F] \rangle + m = 2i,$$

where $[F]$ is the surface in $W_{\mathbf{m}}(K)$ coming from capping off a Seifert surface for K . We can similarly define $\mathfrak{s}_i = \mathfrak{s}_{i_1, \dots, i_\ell}$ as the restriction of Spin^c -structures induced by surgery along links of ℓ components by a similar formula.

We will denote by $\delta(W, \mathfrak{w})$ the quantity

$$\delta(W, \mathfrak{w}) = \frac{c_1(\mathfrak{w})^2 - 3\sigma(W) - 2\chi(W)}{4},$$

which describes the shift in absolute grading induced by the Spin^c -cobordism map on the Floer homology associated to (W, \mathfrak{w}) .

5.1. d -invariants for standard three-manifolds. In this section we review the definition of d -invariants for *standard* three-manifolds from [OS03, Section 9]. An additional reference for this material can be found in [LR14].

Let H be a finitely generated, free abelian group and let $\Lambda^*(H)$ denote the exterior algebra of H . If Y is a three-manifold, we denote

$$\Lambda^* H^1 := \Lambda^* H^1(Y; \mathbb{Z}), \quad \Lambda^* H_1 := \Lambda^*(H_1(Y; \mathbb{Z})/\text{Tors}).$$

The module $HF_{\mathbb{k}}^{\infty}(Y)$ is called *standard* if for each torsion Spin^c structure \mathfrak{t} ,

$$HF_{\mathbb{k}}^{\infty}(Y, \mathfrak{t}) \cong \Lambda^* H^1 \otimes_{\mathbb{Z}} \mathbb{k}[U, U^{-1}]$$

as $\Lambda^* H_1 \otimes_{\mathbb{Z}} \mathbb{k}[U]$ -modules. The group $\Lambda^* H^1$ is graded by setting $\text{gr}(\Lambda^{b_1(Y)} H^1(Y; \mathbb{Z})) = b_1(Y)/2$ and by letting the action of $H_1(Y; \mathbb{Z})/\text{Tors}$ by contraction drop gradings by one.

Let M be any $\Lambda^*(H) \otimes \mathbb{k}$ -module. The kernel of the action of $\Lambda^*(H) \otimes \mathbb{k}$ on M is

$$\mathcal{K}M := \{x \in M \mid v \cdot x = 0 \quad \forall v \in H \otimes \mathbb{k}\}.$$

The quotient of M by this action is defined by

$$\mathcal{Q}M := M/(\mathcal{I} \cdot M).$$

where \mathcal{I} is the two-sided ideal in $\Lambda^*(H) \otimes \mathbb{k}$ generated by H . For a standard three-manifold Y , there are then induced maps:

$$\begin{aligned} \mathcal{K}(\pi) &: \mathcal{K}HF_{\mathbb{k}}^{\infty}(Y, \mathfrak{t}) \rightarrow \mathcal{K}HF_{\mathbb{k}}^+(Y, \mathfrak{t}) \\ \mathcal{Q}(\pi) &: \mathcal{Q}HF_{\mathbb{k}}^{\infty}(Y, \mathfrak{t}) \rightarrow \mathcal{Q}HF_{\mathbb{k}}^+(Y, \mathfrak{t}). \end{aligned}$$

We may now define the *bottom* and *top correction terms* of (Y, \mathfrak{t}) to be the minimal grading of any nonzero element in the image of $\mathcal{K}(\pi)$ and $\mathcal{Q}(\pi)$, denoted by $d_{\text{bot}, \mathbb{k}}$ and $d_{\text{top}, \mathbb{k}}$, respectively.

Proposition 5.1 (Ozsáth-Szabó, [OS03]). *Let $K \subset Y$ be a nullhomologous knot in a three-manifold Y with $b_1(Y) \leq 1$. Then*

$$d_{\text{top}, \mathbb{k}}(Y_0) - \frac{1}{2} \leq d_{\text{top}, \mathbb{k}}(Y_1).$$

Further, if $HF_{\text{red}, \mathbb{k}}(Y) = 0$, then $d_{\text{top}, \mathbb{k}}(Y_0) - \frac{1}{2} = d_{\text{top}, \mathbb{k}}(Y_1)$.

Note that if $b_1(Y) \leq 1$, then $b_1(Y_0) \leq 2$ and $b_1(Y_1) \leq 1$ and hence these manifolds both have standard HF^{∞} [OS03]. Therefore, our use of d_{bot} and d_{top} is justified.

It is natural to ask about three-manifolds which have non-standard HF^{∞} . Links with $\mu_{123} \neq 0$ produce a supply of three-manifolds which are not standard.

Theorem 5.2. *Let \mathcal{L} be an algebraically split link with three components. Suppose that $\mu_{123}(\mathcal{L}) \neq 0$. Then $HF_{\mathbb{Q}}^{\infty}(S_{0,0,0}^3(\mathcal{L}), \mathfrak{s}_0)$ is free of rank 6 over $\mathbb{Q}[U, U^{-1}]$, where \mathfrak{s}_0 is the unique torsion Spin^c structure.*

Proof. Recall that since \mathcal{L} has pairwise linking number zero, there exists a basis a_1, a_2, a_3 for $H^1(S_{0,0,0}^3(\mathcal{L}); \mathbb{Z})$ such that the multiplicity of the triple cup product on cohomology is given by μ_{123} . It now follows immediately from [KM07, Proposition 35.3.2] that $\overline{HM}_{\mathbb{Q}}(S_{0,0,0}^3(\mathcal{L}), \mathfrak{s}_0)$ is free of rank 6 over $\mathbb{Q}[U, U^{-1}]$. Since Heegaard and monopole Floer homology are isomorphic over \mathbb{Z} by [KLT10a, KLT10b, KLT10c, KL11, KL12] or [CGH12b, CGH12c, CGH12a, Tau10], we have the same result for HF^∞ . \square

Remark 5.3. The same proof shows that if \mathbb{k} is an arbitrary field and $\mu_{123}(\mathcal{L})$ is coprime to the characteristic of \mathbb{k} then $HF_{\mathbb{k}}^\infty(S_{0,0,0}^3(\mathcal{L}), \mathfrak{s}_0)$ is free of rank 6 over $\mathbb{k}[U, U^{-1}]$.

5.2. d -invariant inequalities. In this section we collect some inequalities for d -invariants of surgeries of links over an arbitrary field \mathbb{k} . The key result is Proposition 1.14 which shows that the $d_{\mathbb{k}}$ -invariant of $S_1^3(\mathcal{L})$ detects the unlink when \mathcal{L} is an L -space link.

We recall from [OS03] that if (W, \mathfrak{z}) is a negative-definite Spin^c -cobordism from (Y, \mathfrak{s}) to (Y', \mathfrak{s}') , two rational homology spheres, then

$$(14) \quad d(Y', \mathfrak{s}') - d(Y, \mathfrak{s}) \geq \delta(W, \mathfrak{z}).$$

The following three results are well-known consequences of (14) and the formulas for the d -invariants of lens spaces from [OS03]. (Recall that a positive surgery on a nullhomologous knot induces a positive-definite two-handle cobordism; reversing orientation produces a negative-definite cobordism.)

Lemma 5.4. *Let \mathcal{L} be an ℓ -component algebraically split nullhomologous link in a rational homology sphere Y . Fix a Spin^c -structure \mathfrak{s} on Y . For any $\mathbf{m} = (m_1, \dots, m_\ell)$ with $m_i > 0$ we have*

$$(15) \quad d_{\mathbb{k}}(Y_{\mathbf{m}}(L), \mathfrak{s}_{\mathbf{i}}) \leq d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell} d(L(m_k, 1), i_k).$$

where $\mathbf{i} = (i_1, \dots, i_\ell)$.

Corollary 5.5. *Let \mathcal{L} be an algebraically split link in S^3 with ℓ components. Then for any integers $p_1, \dots, p_\ell > 0$ we have*

$$d_{\mathbb{k}}(S_{1/p_1, \dots, 1/p_\ell}^3(\mathcal{L})) \leq d_{\mathbb{k}}(S_1^3(\mathcal{L})).$$

Corollary 5.6. *Let \mathcal{L}' be a sublink in \mathcal{L} . Then*

$$d_{\mathbb{k}}(S_1^3(\mathcal{L})) \leq d_{\mathbb{k}}(S_1^3(\mathcal{L}')).$$

In what follows, it will be important to be able to relate the d -invariants of $+1$ -surgery and large surgery. The key lemma we need is the following.

Lemma 5.7. *Let K be a nullhomologous knot in a rational homology sphere Y with Spin^c structure \mathfrak{s} . Then, for $n \gg 0$, we have*

$$(16) \quad d_{\mathbb{k}}(Y_1(K), \mathfrak{s}_0) \leq d_{\mathbb{k}}(Y_n(K), \mathfrak{s}_0) - d(L(n, 1), 0).$$

Proof. For $n > 0$, consider the cobordism $W : Y \rightarrow Y_1(K)$ given by $W_1(K) \#_{n-1} \mathbb{C}P^2$. Of course, this is a positive-definite four-manifold with diagonalizable intersection form $n\langle 1 \rangle$. We will work with $-W$. Choose a basis $\alpha_1, \dots, \alpha_n$ for $H_2(W)$, where $\alpha_1 = [F_1]$ and the other α_i are given by the exceptional spheres S_i in the $\mathbb{C}P^2$'s. Here, F_1 is a capped off Seifert surface for K . To pin down the signs more carefully, consider the obvious Kirby diagram for $-W$ and handleslide the $n-1$ many -1 -framed unknots onto the -1 -framed copy of $-K$. We now choose the signs on $[F_1]$ and $[S_i]$ so that $[F_n] = \sum_{i=1}^n \alpha_i$ which corresponds to the Seifert surface of the knot K with framing n . (This is pinned down up to an overall sign by the class of $[F_n]$, which will not matter.)

Next, let \mathfrak{z} denote the Spin^c structure on $-W$ which evaluates to one on each of the basis elements. This is the Spin^c structure for which $c_1(\mathfrak{z})^2$ is maximized, i.e. $c_1(\mathfrak{z})^2 = -n$. Note that we can break $-W$ up into two cobordisms $X_1 : -Y \rightarrow -Y_n(K)$ and $X_2 : -Y_n(K) \rightarrow -Y_1(K)$. Of course, each X_i is still negative definite. Let \mathfrak{z}_i denote the restriction of \mathfrak{z} to X_i .

Therefore, we have from (14):

$$(17) \quad d_{\mathbb{k}}(Y_1(K), \mathfrak{s}_0) \leq d_{\mathbb{k}}(Y_n(K), \mathfrak{z}|_{Y_n(K)}) - \delta(X_2, \mathfrak{z}_2).$$

The result will then be complete if we can establish two results. First, we want to see that $\mathfrak{z}|_{Y_n(K)} = \mathfrak{s}_0$. Second, we want to compute that $\delta(X_2, \mathfrak{z}_2) = d(L(n, 1), 0)$. Since F_n is supported in X_1 and $[F_n] = \sum_i \alpha_i$, we see that

$$\langle c_1(\mathfrak{z}_1), [F_n] \rangle = \langle c_1(\mathfrak{z}), [F_n] \rangle = n.$$

The last equality follows since $c_1(\mathfrak{z})$ evaluates to 1 on each α_i . Since $H^2(X_1) = \mathbb{Z}$, \mathfrak{z}_1 is determined by the evaluation of the first Chern class on $[F_n]$. Therefore, it follows that $\mathfrak{z}_1|_{-Y_n(K)} = \mathfrak{s}_0$. (Recall that Spin^c structures do not require an orientation to define, so we can equate the Spin^c structures on $Y_n(K)$ and $-Y_n(K)$.)

Therefore, it remains to compute $\delta(X_2, \mathfrak{z}_2)$. By our choice of \mathfrak{z} ,

$$4\delta(W, \mathfrak{z}) = c_1(\mathfrak{z})^2 - 3\sigma(W) - 2\chi(W) = 0.$$

Next, note that $c_1(\mathfrak{z}_1)^2 - 3\sigma(X_1) - 2\chi(X_1) = -c_1(\mathfrak{z}_2)^2 + 3\sigma(X_2) + 2\chi(X_2)$, since each of these terms is additive over gluing along rational homology spheres. Therefore, we will compute $c_1(\mathfrak{z}_1)^2 - 3\sigma(X_1) - 2\chi(X_1)$ instead. This is easy to compute, since the intersection form of X_1 is $\langle -n \rangle$. Since $c_1(\mathfrak{z}_1)$ is n times the generator of $H^2(X_1)$, we have $c_1(\mathfrak{z}_1)^2 = -\frac{n^2}{n} = -n$. Therefore, we see that

$$\delta(X_1, \mathfrak{z}_1) = \frac{c_1(\mathfrak{z}_1)^2 - 3\sigma(X_1) - 2\chi(X_1)}{4} = \frac{1-n}{4},$$

which is exactly $-d(L(n, 1), 0)$ and we are now done by (17). \square

Corollary 5.8. *For any algebraically split link \mathcal{L} in S^3 we have*

$$(18) \quad d_{\mathbb{k}}(S_1^3(L)) \leq d_{\mathbb{k}}(S_n^3(L), \mathfrak{s}_0) - \sum_{i=1}^{\ell} d(L(n_i, 1), 0),$$

where $\mathbf{n} = (n_1, \dots, n_{\ell})$ is chosen to be a sufficiently large surgery, and \mathfrak{s}_0 denotes the unique trivial Spin^c structure.

The following lemma is a straightforward analogue of [NW15, Theorem 2.5].

Lemma 5.9. *Let K be a nullhomologous knot in a rational homology sphere Y . Choose m a large positive integer and fix a Spin^c structure \mathfrak{s} on Y . Suppose that $d_{\mathbb{k}}(Y_m(K), \mathfrak{s}_0) = d_{\mathbb{k}}(Y, \mathfrak{s}) + d(L(m, 1), 0)$. Then $d_{\mathbb{k}}(Y_m(K), \mathfrak{s}_i) = d_{\mathbb{k}}(Y, \mathfrak{s}) + d(L(m, 1), i)$ for each i .*

Note that $\overline{\mathfrak{s}_i} = \overline{\mathfrak{s}_{-i}}$.

Proof. First, we assume that $0 \leq i \leq \frac{m}{2}$. By the large surgery formula from [OS04a] and absolute gradings on the mapping cone formula [OS11, Section 7.2], $d_{\mathbb{k}}(Y_m(K), \mathfrak{s}_i) = d_{\mathbb{k}}(Y, \mathfrak{s}) + d(L(m, 1), i) - 2H_{\mathfrak{s}}(i)$, where $H_{\mathfrak{s}}(i)$ is defined as follows. The map $v_{\mathfrak{s}, i}^+ : H_*(A_{\mathfrak{s}, i}^+) \rightarrow HF^+(Y, \mathfrak{s})$ defined in [OS08b] is given by multiplication by a power of U when restricted to the image of large powers of U in $H_*(A_{\mathfrak{s}, i}^+)$. This exponent is $H_{\mathfrak{s}}(i)$. Rasmussen shows $H_{\mathfrak{s}}(i) \geq H_{\mathfrak{s}}(i+1) \geq 0$ in [Ras03, Proposition 7.6] and the result follows.

If $i \leq 0$ we use the conjugation symmetry between \mathfrak{s}_i and $\overline{\mathfrak{s}_{-i}}$. □

In fact, we can generalize Lemma 5.9 to links by an induction argument.

Lemma 5.10. *Let \mathcal{L} be a nullhomologous algebraically split ℓ -component link in a rational homology sphere Y . Choose an ℓ -tuple of large positive integers $\mathbf{m} = (m_1, \dots, m_{\ell})$ and fix a Spin^c structure \mathfrak{s} on Y . Suppose*

$$d_{\mathbb{k}}(Y_{\mathbf{m}}(\mathcal{L}), \mathfrak{s}_0) = d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell} d(L(m_k, 1), 0).$$

Then, $d_{\mathbb{k}}(Y_{\mathbf{m}}(\mathcal{L}), \mathfrak{s}_i) = d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell} d(L(m_k, 1), i_k)$ for any tuple $\mathbf{i} = (i_1, \dots, i_{\ell})$.

Proof. We prove this by induction on the number of components in a link in an arbitrary rational homology sphere. If \mathcal{L} is a knot, this is simply Lemma 5.9.

Next, suppose that we have established the result for ℓ -component links in an arbitrary rational homology sphere, and let \mathcal{L} be an $(\ell+1)$ -component link in Y . Let $\mathcal{L}' = \mathcal{L} - L_{\ell+1}$. We will consider two decompositions of the two-handle cobordism $W_{m_1, \dots, m_{\ell+1}}$ from Y to $Y_{\mathbf{m}}(\mathcal{L})$,

$$W_{m_{\ell+1}} \cup X_{m_1, \dots, m_{\ell}} : Y \rightarrow Z \rightarrow Y_{\mathbf{m}}(\mathcal{L}),$$

$$W_{m_1, \dots, m_{\ell}} \cup X_{m_{\ell+1}} : Y \rightarrow Z' \rightarrow Y_{\mathbf{m}}(\mathcal{L})$$

both shown in Figure 4. Here $Z = Y_{m_{\ell+1}}(L_{\ell+1})$ and $Z' = Y_{\mathbf{m}'}(\mathcal{L}')$. Again, the subscripts of the W - and X -labeled cobordisms indicate the components and framings for which the two-handles are attached, as in the notation of the beginning of the section. Let $\overline{\mathcal{L}'}$ be the image of \mathcal{L}' in Z , and $\overline{L_{\ell+1}}$ be the image of $L_{\ell+1}$ in Z' . The Spin^c structures on Y, Z, Z' , and $Y_{\mathbf{m}}(\mathcal{L})$ are denoted as in Figure 4 where $\mathfrak{s}_{i'} = \mathfrak{s}_{i_1, \dots, i_{\ell}}$ and $\mathfrak{s}_{\mathbf{i}} = \mathfrak{s}_{i_1, \dots, i_{\ell+1}}$.




FIGURE 4. Two different decompositions of the two-handle cobordism $W_{m_1, \dots, m_{\ell+1}}$ from Y to $Y_m(\mathcal{L})$ used in Lemma 5.10. The left cobordism consists of first attaching the two-handle to $L_{\ell+1}$ and then attaching the remaining two-handles. The right cobordism consists of first attaching the two-handles along all components other than $L_{\ell+1}$ and then attaching a two-handle along $L_{\ell+1}$

We have

$$\begin{aligned}
 (19) \quad d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell+1} d(L(m_k, 1), 0) &= d_{\mathbb{k}}(Y_m(\mathcal{L}), \mathfrak{s}_0) \\
 &\leq d_{\mathbb{k}}(Z', \mathfrak{s}_{0'}) + d(L(m_{\ell+1}, 1), 0) \\
 &\leq d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell+1} d(L(m_k, 1), 0).
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 (20) \quad d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell+1} d(L(m_k, 1), 0) &= d_{\mathbb{k}}(Y_m(\mathcal{L}), \mathfrak{s}_0) \\
 &\leq d_{\mathbb{k}}(Z, \mathfrak{s}_0) + \sum_{k=1}^{\ell} d(L(m_k, 1), 0) \\
 &\leq d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell+1} d(L(m_k, 1), 0).
 \end{aligned}$$

Combining (19) and (20), we conclude

$$(21) \quad d_{\mathbb{k}}(Z', \mathfrak{s}_{0'}) = d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell} d(L(m_k, 1), 0)$$

$$(22) \quad d_{\mathbb{k}}(Y_m(\mathcal{L}), \mathfrak{s}_0) = d_{\mathbb{k}}(Z, \mathfrak{s}_0) + \sum_{k=1}^{\ell} d(L(m_k, 1), 0).$$

By (20) we get $d_{\mathbb{k}}(Z, \mathfrak{s}_0) = d_{\mathbb{k}}(Y, \mathfrak{s}) + d(L(m_{\ell+1}, 1), 0)$. By applying Lemma 5.9 to the knot $L_{\ell+1}$, we also have

$$(23) \quad d_{\mathbb{k}}(Z, \mathfrak{s}_i) = d_{\mathbb{k}}(Y, \mathfrak{s}) + d(L(m_{\ell+1}, 1), i).$$

Now, using (21), by our induction assumption applied to \mathcal{L}' in Y ,

$$(24) \quad d_{\mathbb{k}}(Z', \mathfrak{s}_{i'}) = d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell} d(L(m_k, 1), i_k).$$

Similarly, by (22), our induction assumption applied to $\overline{\mathcal{L}'}$ in Z implies that

$$(25) \quad d_{\mathbb{k}}(Y_{\mathbf{m}}(L), \mathfrak{s}_{i',0}) = d_{\mathbb{k}}(Z, \mathfrak{s}_0) + \sum_{k=1}^{\ell} d(L(m_k, 1), i_k).$$

Using (23) and (25), we have

$$\begin{aligned} d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell} d(L(m_k, 1), i_k) + d(L(m_{\ell+1}, 1), 0) &\stackrel{(23)}{=} d_{\mathbb{k}}(Z, \mathfrak{t}_0) + \sum_{k=1}^{\ell} d(L(m_k, 1), i_k) \\ &\stackrel{(25)}{=} d_{\mathbb{k}}(Y_{\mathbf{m}}(\mathcal{L}), \mathfrak{s}_{i',0}) \\ &= d_{\mathbb{k}}(Z'_{m_{\ell+1}}(\overline{L_{\ell+1}}), \mathfrak{s}_{i',0}) \\ &\stackrel{(15)}{\leq} d_{\mathbb{k}}(Z', \mathfrak{s}_{i'}) + d(L(m_{\ell+1}, 1), 0) \\ &\stackrel{(24)}{\leq} d_{\mathbb{k}}(Y, \mathfrak{s}) + \sum_{k=1}^{\ell} d(L(m_k, 1), i_k) + d(L(m_{\ell+1}, 1), 0), \end{aligned}$$

hence

$$d_{\mathbb{k}}(Z'_{m_{\ell+1}}(\overline{L_{\ell+1}}), \mathfrak{s}_{i',0}) = d_{\mathbb{k}}(Z', \mathfrak{s}_{i'}) + d(L(m_{\ell+1}, 1), 0).$$

By Lemma 5.9 applied to $\overline{L_{\ell+1}}$ in Z' , we have

$$d_{\mathbb{k}}(Z'_{m_{\ell+1}}(\overline{L_{\ell+1}}), \mathfrak{s}_{i',i_{\ell+1}}) = d_{\mathbb{k}}(Z', \mathfrak{s}_{i'}) + d(L(m_{\ell+1}, 1), i_{\ell+1}).$$

Since $Z'_{m_{\ell+1}}(\overline{L_{\ell+1}}) = Y_{\mathbf{m}}(\mathcal{L})$, (24) now completes the proof. \square

We are now ready to prove the unlink detection theorem for L-space links.

Proof of Proposition 1.14. We prove this by induction on the number of components. First, suppose that \mathcal{L} is a knot. Then, it is well known that $d_{\mathbb{k}}(S_1^3(\mathcal{L})) = 0$ if and only if \mathcal{L} is unknotted.

Next, suppose that the result holds for ℓ -component L-space links and let \mathcal{L} have $\ell + 1$ components. Suppose that $d_{\mathbb{k}}(S_1^3(\mathcal{L})) = 0$. Given $\mathbf{m} \gg \mathbf{0}$, by (18) one has

$$0 = d_{\mathbb{k}}(S_1^3(\mathcal{L})) \leq d_{\mathbb{k}}(S_{\mathbf{m}}^3(\mathcal{L}), \mathfrak{s}_0) - \sum_{k=1}^{\ell+1} d(L(m_k, 1), 0) \leq d(S^3) = 0.$$

The last inequality follows from Lemma 5.4.

By Lemma 5.10, we have that

$$d_{\mathbb{k}}(S_{\mathbf{m}}^3(\mathcal{L}), \mathfrak{s}_i) = \sum_{k=1}^{\ell+1} d(L(m_k, 1), i_k)$$

for all tuples $(i_1, \dots, i_{\ell+1})$. Note that if $d_{\mathbb{k}}(S_1^3(\mathcal{L})) = 0$, then by Lemma 5.6 the same is true for $\mathcal{L}' = L_1 \cup \dots \cup L_{\ell}$. Therefore, we have that \mathcal{L}' is an ℓ -component unlink by the assumption. Hence we have that $S_{\mathbf{m}'}^3(L')$ is $\#_{k=1}^{\ell} L(m_k, 1)$ which is an L-space. Since $\mathbf{m} \gg \mathbf{0}$, we have that $S_{\mathbf{m}}^3(\mathcal{L})$ is an L-space. Therefore, the image of $L_{\ell+1}$ in $S_{\mathbf{m}'}^3(\mathcal{L}')$ is a knot for which $m_{\ell+1}$ -surgery yields a three-manifold with the same absolutely graded Heegaard Floer homology as $S_{\mathbf{m}'}^3(\mathcal{L}') \# L(m_{\ell+1}, 1)$. By Gainullin's Dehn surgery characterization of the unknot in L-spaces [Gai18, Theorem 8.2], we have that $L_{\ell+1}$ is unknotted in $S_{\mathbf{m}'}^3(\mathcal{L}')$. (His paper is only written over \mathbb{F} , but the arguments work over an arbitrary field.) By varying the values of \mathbf{m}' , we may apply [Gab87, Corollary 2.4] to conclude that $L_{\ell+1}$ is actually unknotted in the exterior of \mathcal{L}' . Since \mathcal{L}' is also an unlink, we see that \mathcal{L} is an unlink. The second part directly follows from Corollary 5.5. \square

Remark 5.11. There is a very elementary proof in the case that $\mathbb{k} = \mathbb{F}$. One simply uses (18) to see

$$(26) \quad d_{\mathbb{F}}(S_1^3(\mathcal{L})) \leq d_{\mathbb{F}}(S_{\mathbf{m}}^3(L), \mathfrak{s}_0) - \sum_{k=1}^{\ell} d(L(m_k, 1), 0) = -2h_{\mathcal{L}}(\mathbf{0}) \leq -2.$$

Recall that $-2H_{\mathcal{L}}(\mathbf{0})$ equals $d_{\mathbb{F}}(S_{\mathbf{m}}^3(L), \mathfrak{s}_0)$ up to grading shift which does not depend on the link. Hence one can use the unlink to cancel the grading shift, obtaining the equality in (26). The last inequality follows from the fact that a nontrivial link \mathcal{L} has $h_{\mathcal{L}}(\mathbf{0}) > 0$ [Liu19a].

5.3. A bound from non-vanishing triple linking. In order to constrain the d -invariants of $S_1^3(\mathcal{L})$ in the case that \mathcal{L} is a three-component link with non-trivial Milnor triple linking, we will connect this with the Floer homology of $S_0^3(\mathcal{L})$ which we know is *not* standard by Theorem 5.2. To do this, we will use the d -invariant inequalities (and equalities) that come from the surgery triangle for surgery on nullhomologous knots in three-manifolds which do have standard HF^{∞} . For $n = 0, \infty$ or odd, let \mathfrak{s}_n denote the unique self-conjugate torsion Spin^c structure on $S_{0,0,n}^3(\mathcal{L})$.

Lemma 5.12. *Assume that \mathcal{L} is a Brunnian link with $\mu_{123} \neq 0$. Then*

$$d_{\text{top}, \mathbb{Q}}(S_{0,0,1}^3(\mathcal{L}), \mathfrak{s}_1) \leq d_{\text{bot}, \mathbb{Q}}(S_{0,0,\infty}^3(\mathcal{L}), \mathfrak{s}_{\infty}).$$

Proof. The first step of the proof is to relate the d -invariants of $S_{0,0,\infty}^3(\mathcal{L})$ with $S_{0,0,n}^3(\mathcal{L})$. Choose odd $n \gg 0$. We begin by considering the surgery exact triangle from [OS04b, Theorem 9.19]:

$$\begin{array}{ccc} HF_{\mathbb{Q}}^{\infty}(S_{0,0,\infty}^3(\mathcal{L}), \mathfrak{s}_{\infty}) & \xrightarrow{F_1} & HF_{\mathbb{Q}}^{\infty}(S_{0,0,0}^3(\mathcal{L}), \mathfrak{s}_0) \\ & \swarrow F_3 & \nwarrow F_2 \\ & HF_{\mathbb{Q}}^{\infty}(S_{0,0,n}^3(\mathcal{L}), \mathfrak{s}_n) & \end{array}$$

In the above exact triangle, F_3 is a sum of *two* Spin^c cobordism maps, and by our choice of \mathfrak{s}_n , these have the same absolute grading shift, given by $-d(L(n, 1), 0)$ (See

for instance [OS08b, Section 4.8]). Furthermore, F_3 preserves the absolute $\mathbb{Z}/2$ -grading defined in [OS04b].

Since \mathcal{L} is Brunnian, $S_{0,0,\infty}^3(\mathcal{L})$ is $\#^2(S^1 \times S^2)$. Furthermore, for $Y = S_{0,0,n}^3(\mathcal{L})$ or $S_{0,0,\infty}^3(\mathcal{L})$, since $b_1(Y) \leq 2$, the module

$$(27) \quad HF_{\mathbb{Q}}^{\infty}(Y, \mathfrak{s}) \cong \Lambda^* H^1(Y; \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q}[U, U^{-1}]$$

is standard for any torsion Spin^c structure \mathfrak{s} . The elements of $H_1(Y; \mathbb{Z})/\text{Tors}$ act by contraction on $\Lambda^* H^1(Y; \mathbb{Z})$, hence on $HF_{\mathbb{Q}}^{\infty}(Y, \mathfrak{s})$.

Choose generators x, y of $H_1(S_{0,0,n}^3(\mathcal{L}); \mathbb{Z})/\text{Tors}$. Let us choose a $\mathbb{Q}[U, U^{-1}]$ -basis for $HF_{\mathbb{Q}}^{\infty}(S_{0,0,n}^3(\mathcal{L}), \mathfrak{s}_n)$, denoted $\alpha, \beta, \gamma, \delta$, which correspond to $x^* \wedge y^*, x^*, y^*, 1$ respectively under the isomorphism in (27). Likewise, choose generators x', y' for $H_1(S_{0,0,\infty}^3(\mathcal{L}); \mathbb{Z})/\text{Tors}$ which are bordant to x, y in the surgery cobordism from $S_{0,0,n}^3(\mathcal{L})$ to $S_{0,0,\infty}^3(\mathcal{L})$; define the analogous generators of $HF_{\mathbb{Q}}^{\infty}(S_{0,0,\infty}^3(\mathcal{L}), \mathfrak{s}_{\infty})$ by $\alpha', \beta', \gamma', \delta'$. Then F_3 may be expressed as follows

$$\begin{array}{ccc} \alpha = \langle x^* \wedge y^* \otimes 1 \rangle & \xrightarrow{F_3} & \alpha' = \langle x'^* \wedge y'^* \otimes 1 \rangle \\ \beta = \langle x^* \otimes 1 \rangle \oplus \langle y^* \otimes 1 \rangle = \gamma & & \beta' = \langle x'^* \otimes 1 \rangle \oplus \langle y'^* \otimes 1 \rangle = \gamma' \\ \delta = \langle 1 \otimes 1 \rangle & & \delta' = \langle 1 \otimes 1 \rangle \end{array}$$

By Theorem 5.2 $HF_{\mathbb{Q}}^{\infty}(S_{0,0,0}^3(\mathcal{L}), \mathfrak{s}_0)$ is of rank 6. Exactness implies that

$$HF_{\mathbb{Q}}^{\infty}(S_{0,0,0}^3(\mathcal{L}), \mathfrak{s}_0) \cong \text{coker } F_3 \oplus \ker F_3,$$

and so each of $\ker F_3$ and $\text{coker } F_3$ is rank 3. Recall that the map F_3 is equivariant with respect to the action of the exterior algebra, being a sum of cobordism maps. Hence, the image of α determines the map F_3 . We claim that $F_3(\alpha)$ has a component which is a non-zero multiple of δ' . Before proving the claim, let us see why this will complete the proof. It follows from the long exact sequence relating $HF_{\mathbb{Q}}^{-}, HF_{\mathbb{Q}}^{\infty}$ and $HF_{\mathbb{Q}}^{+}$ that

$$d_{\text{top}, \mathbb{Q}}(S_{0,0,n}^3(\mathcal{L}), \mathfrak{s}_n) - d(L(n, 1), 0) \leq d_{\text{bot}, \mathbb{Q}}(S_{0,0,\infty}^3(\mathcal{L}), \mathfrak{s}_{\infty}).$$

The arguments of Lemma 5.4 establish

$$d_{\text{top}, \mathbb{Q}}(S_{0,0,1}^3(\mathcal{L}), \mathfrak{s}_1) \leq d_{\text{top}, \mathbb{Q}}(S_{0,0,n}^3(\mathcal{L}), \mathfrak{s}_n) - d(L(n, 1), 0),$$

which completes the proof.

Therefore, it remains to prove that $F_3(\alpha)$ contains a non-zero multiple of δ' . Suppose instead that $F_3(\alpha)$ is a non-trivial linear combination of α', β' and γ' .

First, since F_3 is equivariant with respect to the action of the exterior algebra, if $F_3(\alpha) = 0$, then F_3 is identically 0. Thus $\ker F_3$ has rank 4, a contradiction. Since F_3 respects the $\mathbb{Z}/2$ -grading, β' and γ' cannot be components of F_3 . Therefore, it remains to assume $F_3(\alpha) = c\alpha' + d\delta'$ for a unit c (which might involve a nonzero rational factor and a

power of U). Again, since F_3 is equivariant with respect to the action the exterior algebra, contraction by y implies that

$$\begin{aligned} F_3(\beta) &= F_3(x^* \otimes 1) = -F_3 \circ \iota_y(x^* \wedge y^* \otimes 1) \\ &= -\iota_{y'} \circ F_3(x^* \wedge y^* \otimes 1) = -c \cdot \iota_{y'}(x'^* \wedge y'^* \otimes 1) = cx'^* \otimes 1 = c\beta'. \end{aligned}$$

Similarly, $F_3(\gamma) = c\gamma'$ and $F_3(\delta) = c\delta'$. Thus, $\text{rank ker}(F_3) = 0$, which is again a contradiction. \square

Remark 5.13. By Remark 5.3, the above proof works over an arbitrary field \mathbb{k} if $\mu_{123}(\mathcal{L})$ is coprime to the characteristic of \mathbb{k} .

Now we can prove:

Theorem 1.11. *Let $\mathcal{L} = L_1 \cup L_2 \cup L_3$ be an algebraically split link such that all two-component sublinks are \mathbb{Q} -L-space links. If the triple linking number μ_{123} is nonzero, then $d_{\mathbb{Q}}(S_{1,1,1}^3(\mathcal{L})) \leq -2$. If the triple linking number μ_{123} is odd, then the analogous statement holds with \mathbb{Z}_2 -coefficients.*

Proof. Let \mathcal{L} be an algebraically split three-component link such that all two-component sublinks are L-space links. If one of these sublinks is nontrivial, then the result follows from Corollary 5.6 and Proposition 1.14. Therefore from now on we can assume that all two-component sublinks are trivial, so \mathcal{L} is a Brunnian link.

Since $S_{0,0,\infty}^3(\mathcal{L}) = S_{0,0}^3(L_1 \cup L_2) = \#^2(S^1 \times S^2)$, we have $d_{\text{bot},\mathbb{Q}}(S_{0,0,\infty}^3(\mathcal{L})) = -1$. By Lemma 5.12 we get

$$d_{\text{top},\mathbb{Q}}(S_{0,0,1}^3(\mathcal{L})) \leq d_{\text{bot},\mathbb{Q}}(S_{0,0,\infty}^3(\mathcal{L})) = -1.$$

On the other hand, $S_{0,\infty,1}^3(\mathcal{L}) = S^2 \times S^1$ and $S_{\infty,1,1}^3(\mathcal{L}) = S^3$, so by Proposition 5.1 we get

$$d_{\text{top},\mathbb{Q}}(S_{0,0,1}^3(\mathcal{L})) = d_{\text{top},\mathbb{Q}}(S_{0,1,1}^3(\mathcal{L})) + \frac{1}{2} = d_{\mathbb{Q}}(S_{1,1,1}^3(\mathcal{L})) + \frac{1}{2} + \frac{1}{2},$$

and we conclude $d_{\mathbb{Q}}(S_{1,1,1}^3(\mathcal{L})) \leq -2$. A similar argument applies for the case of \mathbb{Z}_2 coefficients by Remark 5.13. (Alternatively, see Corollary 5.20 below.) \square

5.4. 0-surgery on links. In this subsection we describe a different approach to the computation of d -invariants of $S_{000}^3(\mathcal{L})$ building on the work of the second author in [Lid12].¹ Since it uses the link surgery formula of [MO10], we have to restrict ourselves to the coefficients in $\mathbb{F} = \mathbb{Z}_2$.

Recall that the complex $CF^-(S_{000}^3(\mathcal{L}))$ in the unique torsion Spin^c -structure can be written as in Figure 5 using the surgery formula of [MO10].

Theorem 5.14 ([MO10]). *The complex $CF^-(S_{000}^3(\mathcal{L}))$ in the unique torsion Spin^c -structure is quasi-isomorphic (as a complex of free $\mathbb{F}[U]$ -modules) to the complex in Figure 5. The cube filtration on this complex induces a spectral sequence where all pages are link invariants.*

¹As the article appears on arXiv and in thesis form, there is a gap in the argument for $b_1 \geq 5$. This does not affect the arguments used here.




FIGURE 5. Surgery complex quasi-isomorphic to $CF^-(S_{000}^3(\mathcal{L}))$.

After tensoring the surgery complex with $\mathbb{F}[U, U^{-1}]$ over $\mathbb{F}[U]$ and using the fact that $\mathfrak{A}^-(\mathcal{L}, \mathbf{0}) \otimes_{\mathbb{F}[U]} \mathbb{F}[U, U^{-1}]$ is homotopy equivalent to $\mathbb{F}[U, U^{-1}]$ with trivial differential for any link \mathcal{L} , we can simplify the surgery complex for $CF^\infty(S_{000}^3(\mathcal{L}))$. In fact, we can simplify it even further.

Theorem 5.15 ([Lid12]). *The surgery complex for $CF^\infty(S_{000}^3(\mathcal{L}), \mathfrak{s}_0)$ is quasi-isomorphic to the complex in Figure 6. The differentials d_1 and d_2 in the associated spectral sequence vanish, while the relevant d_3 differential is given (up to a unit) by multiplication by the triple linking number $\mu_{123}(\mathcal{L})$ modulo 2.*




FIGURE 6. Surgery complex for computing $HF^\infty(S_{000}^3(\mathcal{L}), \mathfrak{s})$.

Corollary 5.16. $HF^\infty(S_{000}^3(\mathcal{L}), \mathfrak{s}_0)$ has rank 8 if $\mu_{123}(\mathcal{L})$ is even and rank 6 if $\mu_{123}(\mathcal{L})$ is odd.

Recall that the homology of each $\mathfrak{A}^-(\mathcal{L}, \mathfrak{s})$ is non-canonically isomorphic to $\mathbb{F}[U] \oplus M$, where M is annihilated by some power of U . Since U -torsion dies after tensoring with $\mathbb{F}[U, U^{-1}]$, we obtain the following result.

Corollary 5.17. Consider the spectral sequence from Theorem 5.14. The components of the differentials d_1 and d_2 that go between free parts vanish. Up to a power of U , the relevant component of the d_3 differential between free parts is multiplication by $\mu_{123}(\mathcal{L}) \pmod{2}$.

Remark 5.18. Although the differentials d_1 and d_2 between the free parts vanish, the differentials from the free parts to the torsion are still possible, see Example 5.23. Because $H_*(\mathfrak{A}^-(\emptyset)) \cong \mathbb{F}[U]$, there is no torsion piece to which the d_3 differential may map.

We can use these results to give an alternative proof of Theorem 1.11 if the triple linking number is odd and $\mathbb{k} = \mathbb{F}$.

Theorem 5.19. Suppose that $\mu_{123}(L)$ is odd and all two-component sublinks of \mathcal{L} are L -space links. Then $H_{\mathcal{L}}(0, 0, 0) > 0$.

Proof. Let us compute the spectral sequence for $CF^-(S_{000}^3(\mathcal{L}))$. Since all two-component sublinks L_{ij} are L -space links, the components L_i are L -space knots, and

$$H_*(\mathfrak{A}^-(L_{ij}, \mathbf{0})) \cong H_*(\mathfrak{A}^-(L_i), \mathbf{0}) \cong \mathbb{F}[U].$$

Note that the homology of $\mathfrak{A}^-(\mathcal{L}, \mathbf{0})$ might have torsion, since we do not assume \mathcal{L} is an L -space link. By Corollary 5.17 the differentials d_1 and d_2 vanish on the free part of $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{0}))$ and have trivial image in $H_*(\mathfrak{A}^-(\emptyset))$. The differential d_3 from the free part of $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{0}))$ to $H_*(\mathfrak{A}^-(\emptyset)) \cong \mathbb{F}[U]$ is nontrivial.

On the other hand, d_3 lowers the homological degree by 1. Further, up to an absolute shift, the generator of the free part of $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{0}))$ has homological degree $-2H(0, 0, 0)$ while the generator of the free part of $H_*(\mathfrak{A}^-(\emptyset))$ has homological degree -3 , so $-2H(0, 0, 0) - 1 \leq -3$ and $H(0, 0, 0) \geq 1$. \square

Corollary 5.20. Suppose that $\mu_{123}(L)$ is odd and all two-component sublinks of \mathcal{L} are L -space links. Then $d(S_{1,1,1}^3(\mathcal{L})) \leq -2$.

Proof. Similar to Remark 5.11, one simply uses (18) to see that for $\mathbf{m} \gg 0$

$$(28) \quad d(S_{\mathbf{1}}^3(\mathcal{L})) \leq d(S_{\mathbf{m}}^3(L), \mathfrak{s}_0) - \sum_{k=1}^3 d(L(m_k, 1), 0) = -2h_{\mathcal{L}}(\mathbf{0}) = -2H_{\mathcal{L}}(\mathbf{0}) \leq -2.$$

The last inequality follows from Theorem 5.19. \square




FIGURE 7. The link on the left is L_1 from Example 5.21. The link on the right is the result of applying n -twisted Bing doubling to K in L_1 , yielding $L(n) = B(K, n) \cup U$.

5.5. Example: generalized Borromean link. The assumption that all 2-component sublinks are L-space links is important in Theorem 1.11. We will show that there exist three-component algebraically split links \mathcal{L} with nonzero triple linking number and $d(S_{1,1,1}^3(\mathcal{L})) = 0$. Here, we resume working exclusively over \mathbb{F} and omit the coefficients from the notation.

Example 5.21. Start with the two-component link $L_1 = K \cup U$ in the left image of Figure 7 where U is the unknot and K is arbitrary. We can assume the linking number of L_1 is -1 . Let $L(n) = B(K, n) \cup U$ denote the new link obtained by applying an n -twisted Bing-double to K , which is the right image in Figure 7. We order $B(K, n)$ so that the first component is the one “induced” by K . Note that $L(n)$ is a three-component algebraically split link and $\mu_{123}(L(n)) = (-1)\ell k(L_1) = 1$ [Coc90b, Theorem 8.1].

We claim that $d(S_{1,1,1}^3(L(n))) = 0$ for sufficiently large n . Note that $S_{1,1,1}^3(L(n))$ is diffeomorphic to $(S_1^3(D_+(K, n-1)))$ where $D_+(K, n-1)$ is the $(n-1)$ -twisted positively clapped Whitehead double. The d -invariant for this manifold is computed in [Tan17] to be

$$d(S_1^3(D_+(K, n))) = \begin{cases} 0 & n \geq 2\tau(K) \\ -2 & n < 2\tau(K). \end{cases}$$

Hence, for sufficiently large n , we have $\mu_{123}(L(n)) = 1$, but $d(S_{1,1,1}^3(L(n))) = 0$.

Remark 5.22. In the above example,

$$-\lambda(S_{1,1,1}^3(\mathcal{L})) = -\mu_{123}^2(\mathcal{L}) + \beta(B(K, n)) = n - 1.$$

We also have the alternate computation

$$\beta(B(K, n)) = -\lambda(S_{1,1}^3(B(K, n))) = -\lambda(S_1^3(D_+(K, n))) = n.$$

Example 5.23. For a specific example, let K be the unknot. In our conventions $D_+(K, 0)$ is the unknot, $D_+(K, -1) = T(2, 3)$ and $D_+(K, 1)$ is the figure eight knot. In particular, for $n = 0$ we get $L(0)$ is the Borromean rings and

$$d(S_{1,1,1}^3(L(0))) = d(S_1^3(D_+(K, -1))) = -2.$$

The above computation shows that for $n \geq 1$ we have $d(S_{1,1,1}^3(L(n))) = 0$. By a sequence of inequalities similar to (28), we get $H_{L(n)}(0, 0, 0) = 0$ for $n \geq 1$.

For $n = 1$ we can also compute all differentials in the spectral sequence of Theorem 5.14. Indeed, all components of $\mathcal{L} = L(1)$ are unknots and two of three two-component




FIGURE 8. Spectral sequence in Example 5.23 for computing $HF^-(S_{0,0,0}^3(L(1)))$. The solid arrow indicates d_1 while the dashed arrow indicates d_3 . These are the only non-trivial differentials in the spectral sequence after the E_1 page.

sublinks are unlinks. The only interesting two-component sublink is $B(K, 1)$ and in order to apply Theorem 5.14 we need to describe $\mathfrak{A}^-(B(K, 1), \mathbf{0})$. Observe that the trivial component of $B(K, 1)$ has genus 1 in the complement of the other component. This means that for $p \gg 1$ the $(p, 1)$ surgery is large for $B(K, 1)$ and

$$\mathfrak{A}^-(B(K, 1), \mathbf{0}) \simeq CF^-(S_{p,1}^3(B(K, 1), \mathbf{s}_0)) \simeq CF^-(S_p^3(D_+(K, 1), \mathbf{s}_0)) \simeq \mathfrak{A}^-(D_+(K, 1), 0).$$

Here the first and last equations follow from the large surgery formula, and the middle equation is clear.

Since $D_+(K, 1)$ is the figure eight knot, it is well known that $H_*(\mathfrak{A}^-(D_+(K, 1), 0)) \cong \mathbb{F}[U]_{(0)} \oplus \mathbb{F}$. Therefore the E_1 page of the spectral sequence in Theorem 5.14 has the form shown in Figure 8. Let z be the generator of the free part of $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{0}))$. Then, since $H_{\mathcal{L}}(0, 0, 0) = 0$, we see z has degree 3 higher than the generator of $H_*(\mathfrak{A}^-(\emptyset))$. We claim that $d_1(z)$ is the unique non-trivial element in the kernel of U . (This description is independent of the choice of splitting.) Indeed, suppose that instead $d_1(z) = 0$. By Corollary 5.17, d_1 is identically 0, and so is d_2 , and the d_3 differential should map z nontrivially to the homology of $\mathfrak{A}^-(\emptyset)$ which is not possible by degree reasons.

Therefore, $d_1(z)$ is determined and the d_1 differential vanishes elsewhere by Corollary 5.17. On the E_2 page we get a free $\mathbb{F}[U]$ module generated by Uz together with the torsion at the top, and $\mathbb{F}[U]$ everywhere else, so by Corollary 5.17 the differential d_2 vanishes. Now the differential d_3 sends Uz to a power of U times the generator of the homology of $\mathfrak{A}^-(\emptyset)$. For degree reasons, $d_3(Uz)$ is in fact the generator of $H_*(\mathfrak{A}^-(\emptyset))$, and hence this pair of free modules is cancelled by the d_3 differential. The d_3 differential vanishes elsewhere, and all other differentials vanish identically. From the algebra, we cannot seem to determine from the spectral sequence what the torsion coming from

$H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{0}))$ is and whether it contributes to $HF_{red}(S_{0,0,0}^3(\mathcal{L}))$ or the free part. We will use some topological input to complete the spectral sequence computation.

It is an easy Kirby calculus exercise to see that $S_{0,0,0}^3(\mathcal{L}) = \mathbb{T}^3$, and hence $HF_{red}(S_{0,0,0}^3(\mathcal{L})) = 0$. Thus, the torsion term contributes to the free part. Further, ignoring this torsion part, the E_∞ page of the spectral sequence has six towers. Three towers come from the second-to-top filtration level and are all supported in the same gradings. This relies on the fact that $d(S_{p,1}^3(B(K, 1))) = 0$ and that the other two two-component sublinks are trivial. It is also not hard to deduce that this topmost absolute grading is in fact $1/2$. The remaining three towers come from the second-to-bottom filtration level and are all supported in the same gradings; their topmost relative grading is one lower than that of the other towers, and hence have topmost grading $-1/2$. Note that this agrees with the relative-gradings on $HF^-(\mathbb{T}^3)$, and hence the torsion term cannot contribute to the free part of the Floer homology. Consequently, the torsion term of $H_*(\mathfrak{A}^-(\mathcal{L}, \mathbf{0}))$ is trivial, and we have completed the computation of the spectral sequence.

REFERENCES

- [Bak14] Kenneth L. Baker. Counting genus one fibered knots in lens spaces. *Michigan Math. J.*, 63(3):553–569, 2014.
- [BG18] Maciej Borodzik and Eugene Gorsky. Immersed concordances of links and Heegaard Floer homology. *Indiana Univ. Math. J.*, 67(3):1039–1083, 2018.
- [Cas75] A. J. Casson. Link cobordism and Milnor’s invariant. *Bull. London Math. Soc.*, 7:39–40, 1975.
- [CGH12a] Vincent Colin, Paolo Ghiggini, and Ko Honda. The equivalence of Heegaard Floer homology and embedded contact homology III: from hat to plus. arXiv:1208.1526v2 [math.GT], 2012.
- [CGH12b] Vincent Colin, Paolo Ghiggini, and Ko Honda. The equivalence of Heegaard Floer homology and embedded contact homology via open book decompositions I. arXiv:1208.1074v2 [math.GT], 2012.
- [CGH12c] Vincent Colin, Paolo Ghiggini, and Ko Honda. The equivalence of Heegaard Floer homology and embedded contact homology via open book decompositions II. arXiv:1208.1077v2 [math.GT], 2012.
- [CGLS87] Marc Culler, C. McA. Gordon, J. Luecke, and Peter B. Shalen. Dehn surgery on knots. *Ann. of Math. (2)*, 125(2):237–300, 1987.
- [Coc85] Tim D. Cochran. Concordance invariance of coefficients of Conway’s link polynomial. *Invent. Math.*, 82(3):527–541, 1985.
- [Coc90a] Tim D. Cochran. Derivatives of links: Milnor’s concordance invariants and Massey’s products. *Mem. Amer. Math. Soc.*, 84(427):x+73, 1990.
- [Coc90b] Tim D. Cochran. Derivatives of links: Milnor’s concordance invariants and Massey’s products. *Mem. Amer. Math. Soc.*, 84(427):x+73, 1990.
- [Gab87] David Gabai. Foliations and the topology of 3-manifolds. II. *J. Differential Geom.*, 26(3):461–478, 1987.
- [Gai18] Fyodor Gainullin. Heegaard Floer homology and knots determined by their complements. *Algebr. Geom. Topol.*, 18(1):69–109, 2018.
- [Ghi08] Paolo Ghiggini. Knot Floer homology detects genus-one fibred knots. *Amer. J. Math.*, 130(5):1151–1169, 2008.
- [GL89] C. McA. Gordon and J. Luecke. Knots are determined by their complements. *J. Amer. Math. Soc.*, 2(2):371–415, 1989.
- [GLM20] Eugene Gorsky, Beibei Liu, and Allison H Moore. Surgery on links of linking number zero and the Heegaard Floer d -invariant. arXiv:1810.10178, To appear in *Quantum Topology*, 2020.
- [GN15] Eugene Gorsky and András Némethi. Lattice and Heegaard Floer homologies of algebraic links. *Int. Math. Res. Not. IMRN*, (23):12737–12780, 2015.

- [Hos85] Jim Hoste. The first coefficient of the Conway polynomial. *Proc. Amer. Math. Soc.*, 95(2):299–302, 1985.
- [Hos86] Jim Hoste. A formula for Casson’s invariant. *Trans. Amer. Math. Soc.*, 297(2):547–562, 1986.
- [HR85] Craig Hodgson and J. H. Rubinstein. Involutions and isotopies of lens spaces. In *Knot theory and manifolds (Vancouver, B.C., 1983)*, volume 1144 of *Lecture Notes in Math.*, pages 60–96. Springer, Berlin, 1985.
- [KL11] Çağatay Kutluhan and Clifford Henry Lee, Yi-Jen Taubes. HF=HM IV : The Seiberg-Witten Floer homology and ech correspondence. arXiv:1107.2297v1 [math.GT], 2011.
- [KL12] Çağatay Kutluhan and Clifford Henry Lee, Yi-Jen Taubes. HF=HM V : Seiberg-Witten-Floer homology and handle addition. arXiv:1204.0115v3 [math.GT], 2012.
- [KLT10a] Çağatay Kutluhan, Yi-Jen Lee, and Clifford Henry Taubes. HF=HM I : Heegaard Floer homology and Seiberg–Witten Floer homology. arXiv:1007.1979v5 [math.GT], 2010.
- [KLT10b] Çağatay Kutluhan, Yi-Jen Lee, and Clifford Henry Taubes. HF=HM II : Reeb orbits and holomorphic curves for the ech/Heegaard-Floer correspondence. arXiv:1008.1595v2 [math.GT], 2010.
- [KLT10c] Çağatay Kutluhan, Yi-Jen Lee, and Clifford Henry Taubes. HF=HM III : Holomorphic curves and the differential for the ech/Heegaard-Floer correspondence. arXiv:1010.3456v2 [math.SG], 2010.
- [KM07] Peter Kronheimer and Tomasz Mrowka. *Monopoles and three-manifolds*, volume 10 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2007.
- [KMOS07] P. Kronheimer, T. Mrowka, P. Ozsváth, and Z. Szabó. Monopoles and lens space surgeries. *Ann. of Math. (2)*, 165(2):457–546, 2007.
- [KT80] Paik Kee Kim and Jeffrey L. Tolleson. Splitting the PL involutions of nonprime 3-manifolds. *Michigan Math. J.*, 27(3):259–274, 1980.
- [Lid12] Tye Lidman. *Triple Cup Products in Heegaard Floer Homology*. ProQuest LLC, Ann Arbor, MI, 2012. Thesis (Ph.D.)—University of California, Los Angeles.
- [Liu17] Yajing Liu. L-space surgeries on links. *Quantum Topol.*, 8(3):505–570, 2017.
- [Liu19a] Beibei Liu. Four-genera of links and Heegaard Floer homology. *Algebraic & Geometric Topology*, 19(7):3511–3540, 2019.
- [Liu19b] Beibei Liu. Heegaard Floer homology of L-space links with two components. *Pacific J. Math.*, 298(1):83–112, 2019.
- [Liu19c] Beibei Liu. L-space surgeries on 2-component L-space links. *arXiv preprint arXiv:1905.04618*, 2019.
- [LR14] Adam Simon Levine and Daniel Ruberman. Generalized Heegaard Floer correction terms. In *Proceedings of the Gökova Geometry-Topology Conference 2013*, pages 76–96. Gökova Geometry/Topology Conference (GGT), Gökova, 2014.
- [Mil54] John Milnor. Link groups. *Ann. of Math. (2)*, 59:177–195, 1954.
- [Mil57] John Milnor. Isotopy of links. Algebraic geometry and topology. In *A symposium in honor of S. Lefschetz*, pages 280–306. Princeton University Press, Princeton, N. J., 1957.
- [MO10] Ciprian Manolescu and Peter Ozsváth. Heegaard Floer homology and integer surgeries on links. arXiv:1011.1317v4 [math.GT], 2010.
- [Mur66] Kunio Murasugi. On Milnor’s invariant for links. *Trans. Amer. Math. Soc.*, 124:94–110, 1966.
- [Ni09] Yi Ni. Dehn surgeries that yield fibred 3-manifolds. *Math. Ann.*, 344(4):863–876, 2009.
- [NW15] Yi Ni and Zhongtao Wu. Cosmetic surgeries on knots in s_3 . *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2015(706):1–17, 2015.
- [OS03] Peter Ozsváth and Zoltán Szabó. Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary. *Adv. Math.*, 173(2):179–261, 2003.
- [OS04a] Peter Ozsváth and Zoltán Szabó. Holomorphic disks and knot invariants. *Adv. Math.*, 186(1):58–116, 2004.
- [OS04b] Peter Ozsváth and Zoltán Szabó. Holomorphic disks and three-manifold invariants: properties and applications. *Ann. of Math. (2)*, 159(3):1159–1245, 2004.
- [OS05] Peter Ozsváth and Zoltán Szabó. On knot Floer homology and lens space surgeries. *Topology*, 44(6):1281–1300, 2005.

- [OS08a] Peter Ozsváth and Zoltán Szabó. Holomorphic disks, link invariants and the multi-variable Alexander polynomial. *Algebr. Geom. Topol.*, 8(2):615–692, 2008.
- [OS08b] Peter Ozsváth and Zoltán Szabó. Knot Floer homology and integer surgeries. *Algebr. Geom. Topol.*, 8(1):101–153, 2008.
- [OS11] Peter Ozsváth and Zoltán Szabó. Knot Floer homology and rational surgeries. *Algebr. Geom. Topol.*, 11(1):1–68, 2011.
- [OSS15] Peter S. Ozsváth, András I. Stipsicz, and Zoltán Szabó. *Grid homology for knots and links*, volume 208 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [Por80] Richard Porter. Milnor’s $\bar{\mu}$ -invariants and Massey products. *Trans. Amer. Math. Soc.*, 257(1):39–71, 1980.
- [Ras03] Jacob Andrew Rasmussen. *Floer homology and knot complements*. ProQuest LLC, Ann Arbor, MI, 2003. Thesis (Ph.D.)—Harvard University.
- [Smy67] N. Smythe. Isotopy invariants of links and the Alexander matrix. *Amer. J. Math.*, 89:693–704, 1967.
- [Sta65] John Stallings. Homology and central series of groups. *J. Algebra*, 2:170–181, 1965.
- [Sta75] John R. Stallings. Quotients of the powers of the augmentation ideal in a group ring. In *Knots, groups, and 3-manifolds (Papers dedicated to the memory of R. H. Fox)*, pages 101–118. Ann. of Math. Studies, No. 84. 1975.
- [Stu84] Rachel Sturm. *Link concordance invariants (Arf)*. ProQuest LLC, Ann Arbor, MI, 1984. Thesis (Ph.D.)—New York University.
- [Tan17] Motoo Tange. Heegaard Floer homology of Matsumoto’s manifolds. *Adv. Math.*, 320:475–499, 2017.
- [Tau10] Clifford Henry Taubes. Embedded contact homology and Seiberg-Witten Floer cohomology I-V. *Geom. Topol.*, 14(5):2497–3000, 2010.
- [Thu97] William P. Thurston. *Three-dimensional geometry and topology. Vol. 1*, volume 35 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1997. Edited by Silvio Levy.
- [Tor53] Guillermo Torres. On the Alexander polynomial. *Ann. of Math. (2)*, 57:57–89, 1953.
- [Tra84] Lorenzo Traldi. Milnor’s invariants and the completions of link modules. *Trans. Amer. Math. Soc.*, 284(1):401–424, 1984.
- [Tur76] V. G. Turaev. The Milnor invariants and Massey products. *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 66:189–203, 209–210, 1976. Studies in topology, II.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, ONE SHIELDS AVENUE, DAVIS, CA 95616, USA

E-mail address: egorskiy@math.ucdavis.edu

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27607, USA

E-mail address: tlid@math.ncsu.edu

MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY

E-mail address: bbliumath@gmail.com

DEPARTMENT OF MATHEMATICS & APPLIED MATHEMATICS, VIRGINIA COMMONWEALTH UNIVERSITY, 1015 FLOYD AVENUE, BOX 842014, RICHMOND, VA 23284-2014, USA

E-mail address: moorea14@vcu.edu