

BONE MINERAL STUDY

(Tc-99m-MDP, Tc-99m-HMDP)

Overview

- The Bone Mineral Study, with either Tc-99m-MDP or Tc-99m-HMDP, depicts the distribution of bone mineral metabolism throughout the skeleton. In addition, rapid serial images during the first pass of the radiopharmaceutical through the circulation may be obtained to demonstrate regional perfusion.

Indications

- Detection of bone metastases (1-3).
- Diagnosis of osteomyelitis (4,5).
- Evaluation of musculoskeletal trauma (6-9).
- Assessment of low back pain (10).
- Evaluation of primary benign and malignant bone lesions (11).
- Diagnosis of reflex sympathetic dystrophy (12,13).
- Evaluation of the response of Paget's disease to treatment (14).
- Evaluation of heterotopic ossification.

Examination Time

- Initially: 15 minutes for injection of the radiopharmaceutical; 25 minutes for the perfusion and blood pool components of a three phase study.
- 2-3 hours later: 1 hour for image acquisition.

Patient Preparation

- None:
 - > Hydration does not improve image quality (15).
 - > Alendronate does not interfere with radiopharmaceutical uptake (16).

Equipment & Energy Windows

- Gamma camera: Large field of view, preferably with dual heads.
- Collimator: Low energy, high resolution, parallel hole.
- Energy window: 20% window centered at 140 keV.

Radiopharmaceutical, Dose, & Technique of Administration

- Radiopharmaceutical (17,18):
 - Tc-99m-methylene diphosphonate (MDP).

- ❑ Tc-99m-hydroxymethylene diphosphonate (HMDP).
- Dose: 25 mCi (925 MBq).
- Technique of administration:
 - > Routine study: Standard intravenous injection.
 - > Three phase study: Oldendorf method.


Patient Position & Imaging Field

- Patient position: Supine (prone position can be used if patient cannot lie supine).
- Imaging field: Entire body (arms may be omitted).

Acquisition Protocol - Routine Study (17)

- Begin image acquisition 2-3 hours following injection of the radiopharmaceutical.
- Have the patient empty his/her bladder immediately before image acquisition.
- ❑ Moving acquisition protocol:
 1. Use a camera/table motion of approximately 10-15 cm/min.
 2. Acquire ANT and POST images from the head to the bottom to the feet. (If a single head camera is being used, only ANT images of the lower extremities may be acquired.)
 3. Acquire static “spot” images as indicated.
- ❑ Static acquisition protocol:
 1. Acquire an ANT image of the chest for approximately 500-750 K counts, note the time required for acquisition.
 2. Using the same acquisition time used for the ANT chest image acquire ANT and POST images of the rest of the torso and head, and ANT images of the extremities (arms may be omitted).
- SPECT tomographic images are routine for low back pain, and are used in other selected situations (19-22):
 1. Image acquisition parameters:
 - a) degrees of rotation: 360°.
 - b) number of images: 64.
 - c) time per image: 20 seconds.
 2. Data processing:
 - a) reconstruct transverse, sagittal, and coronal image.
 - b) filter selection depends on computer software package.
- Have the patient empty his/her bladder at the end of the study.

Protocol Summary Diagram



Data Processing

- None as a routine unless SPECT imaging is performed.

Optional Maneuvers

- Three phase bone scan (4):
 1. Routinely used when the clinical question is infection in the extremities; frequently used for question of stress fracture, avascular necrosis, or primary bone tumor.
 2. The patient position and field of view depend on the area of interest; include both sides of the body, e.g. both legs or both hands, so that the normal side can be used for comparison.
 3. The radiopharmaceutical is administered as a bolus using the Oldendorf technique:
 - a) if the site of interest is the hands or arms, a tourniquet should be used instead of a blood pressure cuff and there should be a 3 minutes pause between release of the tourniquet and injection (23).
 4. Acquire serial analog images for 5 seconds each for 60 seconds (12 frames) starting at the time of injection.
 5. Immediately acquire a blood pool image for approximately 1 minute. (The number of counts will depend on the body part being imaged and other factors.)
 6. Have the patient return in 2 hours for the delayed images; follow the acquisition protocol for delayed images given above.
- Special views (17):
 1. Images of the scapula with the arms moved forward or up can be used to differentiate activity in the scapula and underlying ribs (24).
 2. The TOD view (tail on detector) is useful for separating otherwise superimposed structures around the pelvic ring. The view is obtained with the patient sitting over the head of the camera.
 3. The skyline view of the elbow improves localization of elbow pathology (25).
 4. Small structures may be magnified with a 2 mm pinhole collimator (26).
- Intraoperative bone imaging: May be used to localize lesions for surgical resection (12,27). The specimen may also be imaged to document that the lesion has been removed.
- Maximum intensity projection (MIP) display: May assist in lesion evaluation (28).
- SPECT-CT imaging: SPECT-CT imaging of the spine is useful in localizing metabolically active vertebral bodies prior to vertebroplasty (29).

Principle Radiation Emission Data - Tc-99m (30)

- Physical half-life = 6.01 hours.

Radiation	Mean % per disintegration	Mean energy (keV)
Gamma-2	89.07	140.5

Dosimetry - Tc-99m-MDP/Tc-99m-HMDP (31,32)

Organ	rads/25 mCi	mGy/925 MBq
Bladder wall		
2 hour void	3.25	32.5
4.8 hour void	7.75	77.5
Kidneys	1.00	10.0
Bone total	0.88	8.8
Red marrow	0.70	7.0
Testes		
2 hour void	0.20	2.0
4.8 hour void	0.28	2.8
Ovaries		
2 hour void	0.30	3.0
4.8 hour void	0.43	4.3
Total body	0.16	1.6

References

- Bombardieri E, Aktolun C, Baum R, et al: Bone scintigraphy: Procedure guidelines for tumour imaging. Eur J Nucl Med Mol Imaging 30:99-106, 2003.
- Krasnow AZ, Hellman RS, Timins ME, et al: Diagnostic bone scanning in oncology. Sem Nucl Med 27:107-141, 1997.
- Terris MK, Klonecke AS, McDougall IR, et al: Utilization of bone scans in conjunction with prostate-specific antigen levels in the surveillance for recurrence of adenocarcinoma after radical prostatectomy. J Nucl Med 32:1713-1717, 1991.
- Holder LE: Clinical radionuclide bone imaging. Radiology 176:607-614, 1990.
- Schauwecker DS: The scintigraphic diagnosis of osteomyelitis. Am J Roentgenol 158:9-18, 1992.
- Karam M, Lavelle WF, Cochiolo J, et al: Vertebral kyphoplasty: Role of skeletal scintigraphy in treatment planning and predicting therapeutic response. J Nucl Med 46:78, 2005.
- Sty JR, Starshak RJ: The role of bone scintigraphy in the evaluation of the suspected abused child. Radiology 146:369-375, 1983.
- Ries T: Detection of osteoporotic sacral fractures with radionuclides. Radiology 146:783-785, 1983.
- Van der Wall H, Storey G, Frater C, et al: Importance of positioning and technical factors in anatomic localization of sporting injuries in scintigraphic imaging. Sem Nucl Med 31:17-27, 2001.
- Connolly LP, d'Hemecourt PA, Connolly SA, et al: Skeletal scintigraphy of young patients with low-back pain and a lumbosacral transitional vertebra. J Nucl Med 44:909-914, 2003.
- Roger B, Bellin MF, Wioland M, et al: Osteoid osteoma: CT-guided percutaneous excision confirmed with immediate follow-up scintigraphy in 16 outpatients. Radiology 201:239-242, 1996.

12. Leitha T, Staudenherz A, Korpan M, et al: Pattern recognition in five-phase bone scintigraphy: Diagnostic patterns of reflex sympathetic dystrophy in adults. Eur J Nucl Med 23:256-262, 1996.
13. O'Donoghue JP, Powe JE, Mattar AG, et al: Three-phase bone scintigraphy: Asymmetric patterns in the upper extremities of asymptomatic normals and reflex sympathetic dystrophy patients. Clin Nucl Med 18:829-836, 1993.
14. Ryan PJ, Gibson T, Fogelman I: Bone scintigraphy following intravenous pamidronate for Paget's disease of bone. J Nucl Med 33:1589-1593, 1992.
15. Klemenz B, Katzwinkel J, Kaiser KP, et al: The influence of differences in hydration on bone-to-soft tissue ratios and image quality in bone scintigraphy. Clin Nucl Med 24:483-487, 1999.
16. Carrasquillo JA, Whatley M, Dyer V, et al: alendronate does not interfere with Tc-99m-methylene diphosphonate bone scanning. J Nuc Med 42:1359-1363, 2001.
17. Kely NL, Cao ZJ, Holder LE: Technical considerations for optimal orthopedic imaging. Sem Nucl Med 27: 328-333, 1997.
18. Littlefield JL, Rudd TG: Tc-99m hydroxymethylene diphosphonate and Tc-99m methylene diphosphonate: Biologic and clinical comparison. J Nucl Med 24:463-466, 1983.
19. Zarnegar K, Siegel ME, Ansari A, et al: Comparison of 180° with 360° SPECT of the spine using single 360° acquisition and MRI confirmation. J Nucl Med 36:122P, 1995.
20. Sarikaya I, Sarikaya A, Holder LE: The role of single photon emission computed tomography in bone imaging. Sem Nucl Med 31:3-16, 2001.
21. Ryan PJ, Evans PA, Gibson T, et al: Chronic low back pain: Comparison of bone SPECT with radiography and CT. Radiology 182:849-854, 1992.
22. Nicholson RL, Mangal-Lan B, Wilkins K: Reverse oblique reconstruction for SPECT of the spine. Clin Nucl Med 16:478-481, 1991.
23. Lecklitner ML, Douglas KP: Increased extremity uptake on three-phase bone scans caused by peripherally induced ischemia prior to injection. J Nucl Med 28:108-111, 1987.
24. Freeman L, Fink-Bennett D: The "hug" view: An image that results in optimal visualization of the upper posterior thorax. Clin Nucl Med 24:63, 1999.
25. Van der Wall H, Magee M, Magnussen J, et al: A novel view for the scintigraphic assessment of the elbow. J Nucl Med 40:194P, 1999.
26. Connolly LP, Treves ST, Davis RT, et al: Pediatric applications of pinhole magnification imaging. J Nucl Med 40:1896-1901, 1999.
27. Wioland M, Serfent-Alaoui A: Didactic review of 175 radionuclide-guided excisions of osteoid osteomas. Eur J Nucl Med 23:1003-1011, 1996.
28. Shih WJ, Magoun S, Stipp V, et al: Volume three-dimensional display of bone SPECT images. J Nucl Med Technol 20:141-146, 1992.
29. Kumar K, Cooins Z, Nguyen M, et al: Incremental benefit of SPECT+CT bone scans over conventional planar and SPECT bone scans in vertebroplasty. J Nucl Med 46:79, 2005.
30. 43-Tc-99m: In MIRD: Radionuclide Data and Decay Schemes, DA Weber, KF Eckerman, AT Dillman, JC Ryman, eds, Society of Nuclear Medicine, New York, 1989, pp 178-179.
31. Graham LS, Krishnamurthy GT, Bland BT: Dosimetry of skeletal-seeking radiopharmaceuticals. J Nucl Med 15:496, 1974.
32. MPI MDP Kit: Product insert. Medi+Physics, 1985.

Normal Findings

- > Weissman BN: Imaging of total hip replacement. Radiology 202:611-623, 1997.
- > Rahmy AIA, Tonino AJ, Tan WD: Quantitative analysis of technetium-99m-methylene diphosphonate uptake in unilateral hydroxyapatite-coated total hip prostheses: First year of follow-up. J Nucl Med 35:1788-1791, 1994.
- > Rubello D, Caricasulo D, Borsato N, Chierichetti F, et al: Three-phase bone scan pattern in asymptomatic uncemented total knee arthroplasty. Eur J Nucl Med 23:1400-1403, 1996.
- > Carnevale V, Dicembrino F, Fruscaiante V, et al: Different patterns of global and regional skeletal uptake of Tc-99m-methylene diphosphonate with age: Relevance to the pathogenesis of bone loss. J Nucl Med 41:1478-1483, 2000.
- > Bar-Sever Z, Connolly LP, Gebhardt MC, et al: Scintigraphy of lower extremity cadaveric bone allografts in osteosarcoma patients. Clin Nucl Med 22:532-535, 1997.
- > Kim HS, Suh JS, Han CD, et al: Sequential Tc-99m MDP bone scans after cementless total hip arthroplasty in asymptomatic patients. Clin Nucl Med 22:6-12, 1997.

[Return to Table of Content](#)